

INDEX

- 16S rDNA, 11–12
- Amplicons, non-specific, 28–29
- Commercial PCR
 - Campylobacter*, 63–64
 - Escherichia coli* O157:H7, 63–64
 - pathogenic, 63–64
 - Listeria monocytogenes*, 63–64
 - Salmonella*, 63–64, 65
- Controls, 28, 32, 46, 59–60
- DNA databases, 11
 - NCBI, 14–19
 - searches, 14–19
- False-negatives, PCR inhibitors, 34–35
- False-positives
 - live vs. dead cells, 33–34
 - non-specific amplicons, 29
 - primer-dimers, 29
 - viable but nonculturable (VBNC), 34
- Interpretation of PCR results, 28–31
- Laboratory
 - equipment, 57–58
 - personnel, 55
 - reagents and disposables, 58–59, 60
 - set-up, 52–55
 - standard operating procedures, 59–60
 - standardization, 62
- Microarrays, 12
- Multiplex PCR, 10, 70, 84
- Multiplex PCR, 70, 84
- Nucleic acid extraction
 - parasites, 120–121
 - viruses, 99–100
- Nucleic acid sequence-based amplification (NASBA), viruses, 102, 107
- parvum*, 124–125
- PCR contamination, 52–55
- PCR design, 12–19
- PCR detection methods, 6–8
 - agarose gels, 6
 - ELISA, 6–8, 56
 - fluorescence, 8–10, 56–57
- PCR detection of
 - Bacillus cereus*, 71
 - Campylobacter*, 63–64, 71–72, 85
 - Clostridium perfringens*, 72–73
 - Cryptosporidium parvum*, 123–124, 144
 - Cyclospora cayetanensis*, 126–127, 144
 - Escherichia coli* O157:H7, 63–64, 85
 - Escherichia coli*, pathogenic, 63–64, 73–76, 84–85
 - Giardia intestinalis*, 128, 145
 - hepatitis A virus, 102, 105–106
 - Listeria monocytogenes*, 63–64, 76–77, 85
 - Microsporidia*, 131–132, 144
 - noroviruses, 100–101, 103–104
 - Salmonella*, 63–64, 85, 77–79, 84–85
 - Shigella*, 79–80
 - Staphylococcus aureus*, 80–81
 - Toxoplasma gondii*, 130–131, 145
 - Vibrio*, pathogenic, 81–83, 85
 - Yersinia enterocolitica*, 83, 85
- PCR inhibitors, 34–35, 46–47
- PCR theory, 2–4
- Quality control, 59–60
- Real-time PCR
 - Campylobacter*, 63–64, 85
 - concept, 8–10
 - Escherichia coli* O157:H7, 63–64, 85
 - pathogenic, 63–64, 85
 - false-positives, 10
 - hepatitis A virus, 108–109
 - Listeria monocytogenes*, 63–64, 85
 - noroviruses, 108–109
 - Salmonella*, 63–64, 85
 - TaqMan, 8
 - Vibrio*, pathogenic, 85,
 - Yersinia enterocolitica*, 85
- Reverse-transcriptase PCR, detection of
 - bacteria, 33–34, 84–85
 - viruses, 101–102
- Sample preparation, 41–49
 - concentrating microbes, 47–49, 93–99, 122–123,
 - dairy, 43, 96–99
 - enrichment, 47–49
 - fruits and produce, 45, 96–99
 - meat, 43–44
 - seafood, 44–45, 94–96
- Sensitivity
 - false-negatives, 31
 - limit of detection, 34–35, 47–49, 70
 - PCR inhibitors, 34–35, 46–47
 - false-positives, 31
- Terminal Restriction Fragment Length Polymorphisms, 10–12

Thermocycler

- conventional, 4, 57–58
- gradient, 5
- hot-air capillary, 4–5, 57–58
- real-time PCR, 5–6, 57–58

Thermocyclers, 4–6, 57–58

Trouble-shooting

- false-negatives, 34–35, 46–47
- false-positives, 32–33, 45

- real-time PCR, 10, 29

- sensitivity, 47–49

Validation of PCR, 31, 60–62

Viability

- bacteria, 33–34

- parasites, 133

Virus, concentration, 93–99