

NAPHTHOQUINONE METABOLITES OF THE FUNGI

A. G. MEDENTSEV and V. K. AKIMENKO*

Institute of Biochemistry and Physiology of Microorganisms, Russian Academy of Sciences, Pushchino, Moscow
Region 142292, Russia

(Received in revised form 25 April 1997)

Key Word Index—Fungi; naphthoquinones; structure; physico-chemical properties; biosynthesis; biogenesis; biological activity; physiological role.

Abstract—Structures and physico-chemical properties of 100 naphthoquinone metabolites produced by filamentous fungi are reviewed. The conditions of pigment formation, biogenesis and the mechanism of biosynthesis of pigments by fungi are described. Sixty-three fungi cultures able to produce naphthoquinone are listed. The biological activities of the main pigments and the mechanism of fungal resistance to their own metabolites are described. The physiological role of the naphthoquinones in producers is discussed. © 1998 Elsevier Science Ltd. All rights reserved

INTRODUCTION

Interaction of species in an ecosystem is known to be determined by many factors, including the ability of the microorganisms to excrete substances which affect the development of other species. Such metabolites called allelotic metabolites play an important role in species adaptation, formation and functioning of communities [1]. The allelotic metabolites cover a broad range of metabolites, including both simple (organic acids, alcohols, sugars etc.) and complex (antibiotics, toxins, hormones etc.) compounds inhibiting and stimulating metabolic processes. This paper presents data on the naphthoquinone secondary metabolites of fungi.

Naphthoquinones are widespread in nature and have been found in higher plants [2], fungi [2, 3] and actinomycetes [4]. The interest of many investigators to this class of compounds is due to their broad-range biological action: phytotoxic [5-8], insecticidal [9], antibacterial [10-17] and fungicidal [12-15]. Besides, some of them also have cytostatic [13] and anticarcinogenic [18-20] properties. Nevertheless, as for most secondary fungal metabolites, at present there is no generally accepted view of the physiological role of naphthoquinones for the producers. In our opinion, it is the high biological activity of the fungal naphthoquinone metabolites with respect to the microorganisms, plants and other species that determines their ecological significance. We consider here the fol-

lowing aspects: (1) structural formulae and available information on the physico-chemical properties of naphthoquinones; (2) biosynthesis and the biogenetic relations of the pigments; and (3) biological activity and the mechanism of action of naphthoquinones. The structures and physico-chemical properties of naphthoquinone metabolites are given in Table 1.

The literature data given in Table 1 indicate that the ability for naphthoquinone biosynthesis is widespread among fungal organisms. It will suffice to note that the list of fungi producers includes 63 cultures (Table 2). Most fungi in this list belong to the genus *Fusarium*. However, it does not mean that other representatives are less active in producing naphthoquinone pigments; they have yet to be examined.

Naphthoquinone metabolites of fungi differ markedly in their chemical composition. Suffice is to compare the simplest of them, juglone, with bicaverine or dimeric naphthoquinones.

CONDITIONS OF NAPHTHOQUINONE BIOSYNTHESIS

The data on the dependence of pigment formation in the fungi on the composition of the medium used have been obtained in studies of naphthoquinone biosynthesis in *Fusarium* [6, 9, 82, 112]. Most of these workers aimed to obtain maximal amounts of various pigments and to study their structure. As a rule, rich complex media were used. Some papers [6, 9, 112] indicate that one of the conditions of naphthoquinone formation is the presence of ammonium nitrogen in the medium. Substitution of nitrate for ammonium

* Author to whom correspondence should be addressed.

Table I. Structures and physico-chemical properties of the Naphthoquinone metabolites of fungi







Naphthoquinone	Same physico-chemical properties
	<p>$C_{11}H_8O_2$ M_r: 172 mp: 90–91° UV λ_{max}^{EtOH} nm (log ϵ): 249 (4.39), 255.5 (4.29), 342.5 (3.44). IR ν^{KBr} cm^{-1}: 1160, 1599. Source: <i>Marasmius graminum</i> [21].</p>
	<p>$C_{10}H_6O_3$ M_r: 174 mp: 164–165° UV λ_{max}^{EtOH} nm (log ϵ): 249 (4.09), 345sh (3.08), 422 (3.56). IR ν^{KBr} cm^{-1}: 1666, 1645, 1603. Source: <i>Verticillium dahliae</i> [22, 23].</p>
	<p>$C_{10}H_6O_4$ M_r: 190.0 UV $\lambda_{max}^{CHCl_3}$ nm (log ϵ): 284 (4.08), 429 (3.54). λ_{max}^{Me-HCl} nm (log ϵ): 282.5 (3.84), 411 (3.36). λ_{max}^{Me-Na} nm (log ϵ): 261 (4.15), 386 (3.20), 470 (3.32). 1H NMR [(CD_3)$_2$CO]: 4.34 (1H), 6.27 (1H), 7.25–7.45 (1H), 7.60–7.81 (2H), 12.70 (1H). Sources: <i>Pyricularia oryzae</i> [24], <i>V. dahliae</i> [25], <i>Cladosporium</i> sp., <i>Stemphylium</i> [26].</p>
	<p>$C_{10}H_6O_4$ M_r: 190.0 UV $\lambda_{max}^{CHCl_3}$ nm (log ϵ): 284 (4.08), 429 (3.54). 1H NMR [(CD_3)$_2$CO]: 4.34 (1H), 6.27 (1H), 7.25–7.45 (1H), 7.60–7.81 (2H), 12.70 (1H). Source: <i>V. dahliae</i> [25].</p>
	<p>$C_{10}H_6O_5$ M_r: 206 mp: 164.5–168.5° UV λ_{max}^{Et-HCl} nm (log ϵ): 213 (4.33), 264 (4.18), 309 (3.86), 399 (2.30), 450 (3.38). UV λ_{max}^{Et-ONa} nm (log ϵ): 284.5 (4.38), 63 (3.78), 554 (3.34). IR ν^{KBr} cm^{-1}: 1592, 1385, 1240, 1175. 1H NMR [(CD_3)$_2$CO]: 6.15 (1H, s), 6.68 (1H, d), 7.15 (1H, d), 12.65 (1H, s). Sources: <i>Aspergillus niger</i> [27], <i>A. citricus</i> [28], <i>Foma vasabiae</i> [23], <i>V. dahliae</i> [25], <i>Cladosporium</i> sp. [26], <i>Stemphylium</i> sp. [26].</p>
	<p>$C_{13}H_{12}O_4$ M_r: 232.0 mp: 103–104° UV λ_{max}^{MeOH} nm (log ϵ): 262 (4.56), 68sh (4.56), 426 (4.01). IR ν^{KBr} cm^{-1}: 2940, 1665, 1635, 1595. 1H NMR ($CDCl_3$): 1.02 (t, Me), 2.64 (q, 2H), 3.85 (s, OMe), 6.73 (s, 2H), 7.05 (s, 1H), 12.10 (s, OH). Source: <i>Aspergillus parvulus</i> [29].</p>


Table 1—continued.

Naphthoquinone	Same physico-chemical properties
	<p>$C_{11}H_8O_6$, M_r: 236 mp: 255 UV λ_{max}^{EtOH} nm (log ϵ): 225 (4.10), 269.5 (4.1), 320 (3.82), 384 (3.50), 470sh (3.08). UV λ_{max}^{EtONa} nm: 227, 291, 374, 580. IR ν^{KBr} cm^{-1}: 3360, 1650, 1615. 1H NMR (DMSO-d_6): 3.88 (s, OMe), 6.50 (s, H-Ar), 6.94 (s, H-Ar), 12.20 (s, OH). Source: <i>Cercospora melonis</i> [30].</p>
(7) 2-methoxy-3,6,8-trihydroxy-1,4-NQ	
	<p>$C_{13}H_{12}O_4$, M_r: 232 mp: 116–117° UV λ_{max}^{EtOH} nm (log ϵ): 217 (4.0), 287 (4.17), 410 (2.61). IR ν^{KBr} cm^{-1}: 3360, 1660, 1625, 1615. Source: <i>Foma vasabiae</i> [31].</p>
(8) 2-hydroxy-6-methoxy-3,5-dimethyl-1,4-NQ	
	<p>$C_{11}H_8O_6$, M_r: 236 mp: 267° UV λ_{max}^{EtOH} nm (log ϵ): 272 (4.33), 285sh, 322 (3.99), 385 (3.56), 485 (3.26). IR ν^{KBr} cm^{-1}: 3480, 1660, 1620. Source: <i>Corynespora casiocala</i> [32].</p>
(9) 7-methylspinochrome B	
	<p>$C_{15}H_{16}O_7$, M_r: 308 mp: 202 UV λ_{max}^{EtOH} nm (log ϵ): 273 (4.26), 286sh, 331 (3.87), 400 (3.44). IR ν^{KBr} cm^{-1}: 3400, 1650, 1610. Source: <i>Corinespora casiocala</i> [32].</p>
(10) 7-(3-hydroxy-n-butyl)-6-methyl-spinochrome B	
	<p>$C_{14}H_{14}O_6$, M_r: 262 mp: 187° UV λ_{max}^{MeOH} nm: 221, 256, 262, 305, 422. IR ν^{KBr} cm^{-1}: 1672, 1628, 1590. 1H NMR (CDCl$_3$): 1.13 (t, Me), 2.75 (q, CH$_2$), 3.92 (s, OMe), 4.00 (s, OMe), 6.04 (s, H-Ar), 7.30 (s, H-Ar), 12.50 (s, OH). Source: <i>Hendersonula toruloidea</i> [33].</p>
(11) 2,7-dimethoxy-5-hydroxy-6-ethyl-1,4-NQ	
	<p>$C_{16}H_{14}O_7$, M_r: 320 mp: 160–163° UV λ_{max}^{MeOH} nm (log ϵ): 220 (4.51), 57sh (4.14), 263 (4.15), 306 (3.97), 425 (3.55). IR ν^{CCl_4} cm^{-1}: 1742, 1690, 1632. Source: <i>Hendersonula toruloidea</i> [34].</p>
(12) 2,7-dimethoxy-6-(acetoxylethyl)juglone	
	<p>$C_{10}H_6O_6$, M_r: 222 mp: >300°-decomp. UV λ_{max}^{EtOH} nm (log ϵ): 228 (4.43), 272 (4.06), 318 (3.93), 486 (3.75), 517 (3.80). IR ν^{KBr} cm^{-1}: 3240, 1660, 1625, 1602. 1H NMR (DMSO-d_6): 6.31 (s, H-Ar), 6.31 (s, H-Ar), 13.1 (s, 2OH). Source: <i>Helicobasidium mompa</i> [35].</p>
(13) Mompain	

Table 1—continued.


Naphthoquinone

Same physico-chemical properties




(14) 2,7-dimethylnaphthazarin

$C_{12}H_{10}O_4$ M_r : 194 mp: 125–126
 UV λ_{max}^{EtOH} nm (log ϵ): 217 (4.55), 280 (3.30), 480 (3.77), 510 (3.79),
 550 (3.58).
 IR ν^{KBr} cm^{-1} : 1611.
 Source: *Mollisia caesia* [36].




(15) Mollisin

$C_{14}H_{10}O_4Cl_2$ M_r : 313 mp: 202–203°
 UV λ_{max}^{EtOH} nm (log ϵ): 259 (4.26), 280 (3.90), 420 (3.52).
 IR ν^{KBr} cm^{-1} : 1720, 1649, 1618.
 1H NMR ($CDCl_3$) (τ): -2.05 (OH), 2.81 (H-Ar), 3.67 (COCHCl₂),
 7.58 (CH-Ar), 7.84 (CH-Q), 2.81 (H-Q).
 Sources: *Mollisia caesia*, *M. fallens* [36–38].




(16) 2,5,7-trihydroxy-3-(5-hydroxyhexyl)-1,4-NQ

$C_{16}H_{18}O_6$ M_r : 306 mp: 166–168°
 UV λ_{max}^{EtOH} nm (log ϵ): 214, 263, 314, 382, 460.
 IR ν^{KBr} cm^{-1} : 3370, 1650–1630, 1340.
 1H NMR (acetone- d_6): 1.09 (s, Me), 1.04 (2Me), 2.54 (2H), 3.70
 (1H), 6.59 (H-Ar), 7.06 (H-Ar), 12.68 (OH).
 Source: *Penicillium* sp. [39].




(17) Trichione

$C_{17}H_{14}O_6$ M_r : 346
 Source: *Trichia floriformis* [40].




(18) Homotrichione

$C_{19}H_{18}O_8$ M_r : 374
 Source: *Metatrichia vespasarium* [40].



(19) 2-acetyl-3,5-dihydroxy-7-methoxy-1,4-NQ

$C_{14}H_{12}O_6$ M_r : 276 mp: 208°
 UV λ_{max}^{MeOH} nm (log ϵ): 208 (4.4), 227 (4.3), 272 (4.2), 313 (4.1), 380
 (3.7).
 IR ν^{KCl} cm^{-1} : 3340, 2980, 1705, 1635, 1610.
 1H NMR (pyridine- d_5): 2.30 (s, Me), 3.70 (s, OMe), 3.95 (s, CH₂),
 6.70 (d, H-Ar), 7.35 (d, H-Ar), 10.7 (s, OH).
 Source: *Cylindrocarpon* sp. [41].



(20) 1,4,5,8-tetrahydroxy-3-ethylnaphthalene-2,6-dione

$C_{12}H_{10}O_6$ M_r : 250 mp: 183–185°
 UV λ_{max}^{MeOH} nm (log ϵ): 217, 241sh, 266, 485sh, 517, 555.
 IR ν^{KBr} cm^{-1} : 3527, 3400, 2968–2872, 1598, 1460.
 1H NMR ($CDCl_3$): 1.23 (t, 3H, Me), 2.66 (q, 2H, CH₂), 11.33 (s, OH).
 Source: *Cetraria cucullata* [42].

Table 1—continued.









Naphthoquinone	Same physico-chemical properties
 <p data-bbox="41 439 220 459">(21) Homoventosin</p>	<p data-bbox="556 368 884 389">$C_{15}H_{14}O_6$, M_r: 306 mp: 180° decomp.</p> <p data-bbox="556 391 1049 411">UV λ_{max}^{EtOH} nm (log ϵ): 235 (4.43), 294 (3.98), 505 (3.90).</p> <p data-bbox="556 413 897 433">IR ν^{KBr} cm^{-1}: 1725, 1662, 1610, 1570.</p> <p data-bbox="556 435 888 455">Source: <i>Haematomma ventosum</i> [43].</p>
 <p data-bbox="41 681 166 701">(22) Herbarin</p>	<p data-bbox="556 560 841 580">$C_{16}H_{16}O_6$, M_r: 304 mp: 192–193°</p> <p data-bbox="556 582 1009 602">IR ν^{KBr} cm^{-1}: 3325, 1678, 1650, 1607, 1572, 1567.</p> <p data-bbox="556 604 1171 675">1H NMR (pyridine-d_5) (τ): 2.61 (<i>d</i>, 1H), 3.18 (<i>d</i>, 1H), 4.9–5.0 (<i>m</i>, 2H), 5.4 (<i>s</i>, 1H), 6.17 (<i>s</i>, 3H), 6.21 (<i>s</i>, 3H), 6.86 (<i>m</i>, 1H), 7.33 (<i>m</i>, 1H), 8.26 (<i>s</i>, 3H).</p> <p data-bbox="556 677 825 697">Source: <i>Torula herbarum</i> [44].</p>
 <p data-bbox="41 915 243 935">(23) Dehydroherbarin</p>	<p data-bbox="556 772 841 792">$C_{16}H_{14}O_6$, M_r: 286 mp: 189–190°</p> <p data-bbox="556 794 1171 844">UV λ_{max}^{EtOH} nm (log ϵ): 217 (4.12), 250 (3.85), 272 (3.85), 335 (3.28), 400 (3.15), 485 (3.15).</p> <p data-bbox="556 846 951 866">IR ν^{KBr} cm^{-1}: 1675, 1632, 1610, 1600, 1565.</p> <p data-bbox="556 868 1171 919">1H NMR ($CDCl_3$) (τ): 2.75 (1H, <i>d</i>), 3.30 (1H, <i>d</i>), 4.18 (1H, <i>d</i>), 4.88 (2H, <i>s</i>), 6.05 (3H, <i>s</i>), 6.07 (3H, <i>s</i>), 8.02 (3H, <i>s</i>).</p> <p data-bbox="556 921 825 941">Source: <i>Torula herbarum</i> [44].</p>
 <p data-bbox="41 1157 247 1177">(24) O-methylherbarin</p>	<p data-bbox="556 1054 841 1074">$C_{17}H_{18}O_6$, M_r: 318 mp: 188–190°</p> <p data-bbox="556 1076 874 1096">UV λ_{max}^{EtOH} nm: 218, 267, 285sh, 415.</p> <p data-bbox="556 1098 1171 1149">1H NMR ($CDCl_3$) (τ): 2.73 (H-Ar), 5.25–5.65 (2H), 7.2–7.55 (2H), 6.06–6.08 (2OMe-Ar), 6.72 (OMe), 8.52 (Me).</p> <p data-bbox="556 1151 825 1171">Source: <i>Torula herbarum</i> [45].</p>
 <p data-bbox="41 1399 198 1419">(25) Lambertellin</p>	<p data-bbox="556 1296 841 1316">$C_{14}H_8O_5$, M_r: 256 mp: 253–254°</p> <p data-bbox="556 1318 1049 1338">UV λ_{max}^{EtOH} nm (log ϵ): 284 (4.08), 290 (4.10), 430 (3.68).</p> <p data-bbox="556 1340 905 1360">IR ν^{NaCl} cm^{-1}: 1746, 1665, 1657, 1620.</p> <p data-bbox="556 1362 1171 1382">1H NMR (D_2O) (τ): 1.74–2.58 (<i>m</i>, Me), 3.23 (<i>q</i>, 1H), 7.29 (<i>d</i>, 3H).</p> <p data-bbox="556 1384 1085 1405">Sources: <i>Lambertella</i> sp. [46], <i>Pseudospirones simplex</i> [47].</p>
 <p data-bbox="41 1641 166 1661">(26) Gunacin</p>	<p data-bbox="556 1497 713 1518">$C_{17}H_{16}O_6$, M_r: 348</p> <p data-bbox="556 1520 942 1540">UV λ_{max}^{EtOH} nm: 228, 267, 311–317, 430–445.</p> <p data-bbox="556 1542 905 1562">λ_{max}^{EtONa} nm: 238, 290, 310–320, 520–545.</p> <p data-bbox="556 1564 897 1584">IR ν^{KBr} cm^{-1}: 1730–1740, 1635, 1250.</p> <p data-bbox="556 1586 1171 1636">1H NMR ($CDCl_3$): 7.17 (1H, <i>d</i>), 6.65 (1H, <i>d</i>), 6.2 (1H, <i>d</i>), 4.4 (1H, <i>m</i>), 4.15 (1H, <i>m</i>), 3.9 (3H, <i>s</i>), 2.1 (3H, <i>s</i>), 1.7 (3H, <i>d</i>).</p> <p data-bbox="556 1639 780 1659">Source: <i>Ustilago</i> sp. [15].</p>
 <p data-bbox="41 1882 206 1903">(27) Cryptosporin</p>	<p data-bbox="556 1780 794 1800">$C_{14}H_{12}O_6$, M_r: 276 mp: 244</p> <p data-bbox="556 1802 1049 1822">UV λ_{max}^{MeOH} nm (log ϵ): 240 (4.08), 287 (4.00), 408 (3.65).</p> <p data-bbox="556 1824 951 1844">IR ν^{KBr} cm^{-1}: 3550, 3500, 3300, 1660, 1635.</p> <p data-bbox="556 1846 884 1866">Source: <i>Cryptosporium pinicola</i> [48].</p>

Table 1—continued.


Naphthoquinone

Same physico-chemical properties




(28) Rodocladonic acid

$C_{15}H_{10}O_8$ M_r : 318 mp: $>300^\circ$ decomp.
 UV λ_{max}^{EtOH} nm (log ϵ): 294 (4.37), 318 (4.18), 328 (3.99), 449 (3.65).
 IR ν^{KBr} cm^{-1} : 3385, 3300, 1660, 1630.
 Source: *Cladonia* sp. (lichens) [49].




(29) Kanarion

$C_{14}H_8O_8$ M_r : 272 mp: 192–197°
 UV λ_{max}^{MeONa} nm (log ϵ): 241 (4.15), 267 (4.03), 370 (3.43), 430 (3.40).
 λ_{max}^{MeONa} nm: 279, 384, 520.
 IR ν^{KBr} cm^{-1} : 3450, 2940, 2850, 1650, 1595.
 Sources: *Usnea canariensis*, *Usnea nookeri* (lichens) [50].




(30) 6-methyl-dehydro-a-lapachone

$C_{16}H_{14}O_8$ M_r : 254 mp: 130–132°
 UV λ_{max}^{EtOH} nm (log ϵ): 204 (4.52), 265 (4.30), 275 (4.26), 333 (3.76), 444 (3.14).
 IR ν^{KBr} cm^{-1} : 1668, 1648.
 1H NMR ($CDCl_3$): 1.57 (s, 2Me), 2.51 (s, Me), 5.75 (d, 1H), 6.70 (d, 1H), 7.93 (s, 1H), 7.55 (d, 1H), 8.02 (d, 1H).
 Source: *Fomes annosus* [51].




(31) Phomasarin

$C_{19}H_{16}O_8N$ M_r : 387 mp: 196°
 UV λ_{max}^{EtOH} nm (log ϵ): 231 (4.54), 277 (4.77), 430 (3.92).
 IR ν^{CHCl_3} cm^{-1} : 1633, 1694.
 1H NMR (CF_3COOD) (τ): 1.95 (s, 1H), 5.75 (s, OMe), 7.06 (t, 2H, CH_2), 8.4 (m, 4H, 2 CH_2), -2.48 (s, OH), 8.95 (t, Me).
 Source: *Pyrenochaeta cerestris* [52].




(32) Isophomasarin

$C_{18}H_{17}O_8N$ M_r : 387 mp: 215–216°
 UV λ_{max}^{EtOH} nm (log ϵ): 231 (4.54), 277 (4.77), 430 (3.92).
 IR ν^{CHCl_3} cm^{-1} : 1615, 1730.
 1H NMR (CF_3COOD): 1.72 (s, 1H), 5.86 (s, OMe), 7.06 (2H, CH_2), 8.4 (4H, 2 CH_2), 8.97 (t, Me).
 Source: *Pyrenochaeta cerestris* [53].



(33) Erythrostrominone

$C_{17}H_{16}O_8$ M_r : 348 mp: 184–185°
 UV λ_{max}^{EtOH} nm (log ϵ): 231.5 (4.54), 280 (3.89), 315 (3.9), 480 (3.86), 509 (3.93), 546 (3.72).
 IR ν^{CHBr_3} cm^{-1} : 3580, 1720, 1604.
 1H NMR ($CDCl_3$) (τ): 5.15 (m, 1H), 5.38 (m, 1H), 8.25 (m, 2H), 3.73 (s, H-Ar), 7.1 (s, 2H), 7.77 (s, 3H), 6.16 (s, OMe), -2.6 (s, OH), -3.9 (s, OH).
 Source: *Gnomonia erythrostroma* [54].



(34) Deoxyerythrostrominone

$C_{17}H_{16}O_7$ M_r : 332 mp: 148–150°
 UV λ_{max}^{EtOH} nm (log ϵ): 234 (4.49), 275 (3.83), 317 (3.88), 477 (3.84), 505 (3.91), 542 (3.73).
 IR ν^{CHBr_3} cm^{-1} : 1713, 1601.
 1H NMR ($CDCl_3$) (τ): 5.47 (m, 1H), 8.3 (m, 2H), 7.4 (m, 2H), 7.1 (m, 2H), 7.74 (s, Me), 3.72 (s, H-Ar), 6.16 (s, Me), -2.6 (s, OH), -3.9 (s, OH).
 Source: *Gnomonia erythrostroma* [54].

Table I—continued.








Naphthoquinone	Same physico-chemical properties
 <p data-bbox="45 431 297 459">(35) Deoxyerythrostominol</p>	<p data-bbox="561 268 848 292">$C_{17}H_{18}O_7$, M_r: 334 mp: 139–141°</p> <p data-bbox="561 294 1176 338">UV λ_{max}^{EtOH} nm (log ϵ): 234 (4.46), 275 (3.85), 317 (3.89), 475sh (3.84), 505 (3.92), 541 (3.73).</p> <p data-bbox="561 340 801 364">IR ν^{Nujol} cm^{-1}: 3580, 1603.</p> <p data-bbox="561 366 1176 431">1H NMR ($CDCl_3$) (τ): 5.59 (<i>m</i>, 1H), 8.12 (<i>m</i>, 2H), 7.37 (<i>m</i>, 1H), 3.65 (<i>s</i>, H-Ar), 7.1 (<i>m</i>, 2H), 5.82 (<i>m</i>, 1H), 8.72 (3H), 6.16 (<i>s</i>, OMe). —2.6 (<i>s</i>, OH), —3.9 (OH).</p> <p data-bbox="561 433 895 457">Source: <i>Gnomonia erythrostoma</i> [54].</p>
 <p data-bbox="45 661 270 689">(36) Epierythrostominol</p>	<p data-bbox="561 497 848 522">$C_{17}H_{18}O_8$, M_r: 350 mp: 187–191°</p> <p data-bbox="561 524 1176 568">UV λ_{max}^{EtOH} nm (log ϵ): 231 (4.51), 277 (3.91), 317 (3.90), 482 (3.84), 514 (3.93), 551 (3.73).</p> <p data-bbox="561 570 801 594">IR ν^{Nujol} cm^{-1}: 3570, 1605.</p> <p data-bbox="561 596 1176 661">1H NMR ($CDCl_3$) (τ): 8.67 (<i>d</i>, Me), 7.97 (<i>m</i>, 2CH_2), 6.05 (<i>s</i>, OMe), 5.72 (<i>m</i>, 2H), 5.04 (<i>m</i>, 1H), 3.58 (<i>s</i>, 1H), —2.63 (<i>s</i>, OH), —3.21 (<i>s</i>, OH).</p> <p data-bbox="561 663 895 687">Source: <i>Gnomonia erythrostoma</i> [55].</p>
 <p data-bbox="45 911 182 939">(37) Fusarubin</p>	<p data-bbox="561 719 801 743">$C_{15}H_{14}O_7$, M_r: 306 mp: 218</p> <p data-bbox="561 745 901 770">UV λ_{max}^{EtOH} nm: 224, 304, 475, 500, 535.</p> <p data-bbox="561 772 1022 796">IR ν^{CDCl_3} cm^{-1}: 3550, 3425, 2900, 2825, 1600, 1570.</p> <p data-bbox="561 798 1176 862">1H NMR ($CDCl_3$): 1.80 (3H, <i>s</i>, Me), 3.05 (2H, <i>m</i>), 3.80 (3H, <i>s</i>, OMe), 5.15 (2H, <i>m</i>), 6.33 (1H, <i>s</i>, H-Ar), 12.69 (1H, <i>s</i>, OH), 12.95 (1H, <i>s</i>, OH).</p> <p data-bbox="561 864 1176 929">Sources: <i>Fusarium solani</i> [7, 11, 58, 59], <i>Neocosmospora vasinfectum</i>, <i>N. africana</i> [57], <i>Nectria haematococca</i> [56], <i>F. decemcellulare</i> [60], <i>F. moniliforme</i> [61].</p>
 <p data-bbox="45 1145 236 1173">(38) Deoxyfusarubin</p>	<p data-bbox="561 1001 848 1026">$C_{15}H_{14}O_6$, M_r: 290 mp: 208–212°</p> <p data-bbox="561 1028 1176 1072">UV λ_{max}^{EtOH} nm (log ϵ): 218 (3.95), 250 (3.84), 295 (3.72), 410 (3.33), 423 (3.34), 445 (3.22).</p> <p data-bbox="561 1074 790 1098">IR ν^{KBr} cm^{-1}: 1680, 1620.</p> <p data-bbox="561 1100 1176 1145">1H NMR ($CDCl_3$): 1.65 (3H, <i>s</i>), 2.96 (2H, <i>s</i>), 3.94 (3H, <i>s</i>), 4.92 (2H, <i>s</i>), 6.04 (1H, <i>s</i>), 7.43 (1H, <i>s</i>), 12.25 (1H, <i>s</i>).</p> <p data-bbox="561 1147 874 1171">Source: <i>Nectria haematococca</i> [62].</p>
 <p data-bbox="45 1378 260 1407">(39) O-methylfusarubin</p>	<p data-bbox="561 1235 848 1260">$C_{16}H_{16}O_7$, M_r: 320 mp: 138–139°</p> <p data-bbox="561 1262 1176 1306">UV λ_{max}^{MeOH} nm (log ϵ): 226 (4.48), 282.5 (4.05), 484 (3.83), 510 (3.80), 550 (3.49).</p> <p data-bbox="561 1308 989 1332">IR ν^{KBr} cm^{-1}: 3470, 3240, 1605, 1550, 1460 [63].</p> <p data-bbox="561 1334 1176 1378">1H NMR ($CDCl_3$): 1.56 (3H, Me), 2.50–2.78 (2H), 4.01 (3H, OMe), 4.05 (3H, OMe), 4.66 (2H), 6.85 (1H, H-Ar), 13.16 (1H, OH).</p> <p data-bbox="561 1380 1056 1405">Sources: <i>Fusarium moniliforme</i> [61], <i>F. oxysporum</i> [63].</p>
 <p data-bbox="45 1612 283 1641">(40) O-demethylfusarubin</p>	<p data-bbox="561 1489 848 1514">$C_{14}H_{12}O_7$, M_r: 292 mp: 195–197°</p> <p data-bbox="561 1516 901 1540">UV λ_{max}^{EtOH} nm: 230, 305, 475, 505, 540.</p> <p data-bbox="561 1542 897 1566">IR ν^{KBr} cm^{-1}: 3450, 3380, 1620, 1575.</p> <p data-bbox="561 1568 1176 1612">1H NMR ($CDCl_3$): 1.65 (3H, Me), 2.67–2.96 (2H), 4.85–4.91 (2H), 6.34 (1H, H-Ar), 13.01 (1H, OH), 11.88 (1H, OH).</p> <p data-bbox="561 1614 830 1639">Source: <i>F. decemcellulare</i> [64].</p>
 <p data-bbox="45 1846 260 1874">(41) O-methylfusarubin</p>	<p data-bbox="561 1683 848 1707">$C_{16}H_{16}O_8$, M_r: 320 mp: 188–190°</p> <p data-bbox="561 1709 1176 1753">UV λ_{max}^{EtOH} nm: 225 (4.49), 302 (3.95), 470 (3.78), 495 (3.83), 532 (3.64).</p> <p data-bbox="561 1755 790 1780">IR ν^{KBr} cm^{-1}: 3425, 1600.</p> <p data-bbox="561 1782 1176 1846">1H NMR ($CDCl_3$): 1.54 (3H, <i>s</i>, Me), 3.04–2.64 (2H, <i>dd</i>), 3.31 (3H, <i>s</i>, OMe), 3.92 (3H, <i>s</i>, OMe), 4.90–4.53 (2H, <i>dd</i>), 6.17 (1H, <i>s</i>, H-Ar), 12.63 (1H, <i>s</i>, OH), 12.91 (1H, <i>s</i>, OH).</p> <p data-bbox="561 1848 888 1872">Sources: <i>F. martii</i> [13], <i>F. solani</i> [65].</p>

Table 1—continued.








Naphthoquinone	Same physico-chemical properties
 <p data-bbox="135 439 333 465">(42) <i>O</i>-ethylfusarubin</p>	<p data-bbox="650 298 935 324">$C_{17}H_{18}O_7$ M_r: 334 mp: 138–139</p> <p data-bbox="650 324 1143 350">UV λ_{max}^{EtOH} nm (log ϵ): 225 (4.44), 302 (3.91), 470 (3.78).</p> <p data-bbox="650 350 884 376">IR ν^{KBr} cm^{-1}: 3425, 1625.</p> <p data-bbox="650 376 1267 439">1H NMR ($CDCl_3$): 1.55 (3H, <i>s</i>, Me), 3.04–2.64 (2H, <i>dd</i>), 3.61–1.55 (5H, <i>q</i>, <i>t</i>, OCH_2CH_3), 3.92 (3H, <i>s</i>, OMe), 4.88–4.52 (2H, <i>dd</i>), 6.17 (1H, <i>s</i>, H-Ar), 12.63 (1H, <i>s</i>, OH), 12.91 (1H, <i>s</i>, OH).</p> <p data-bbox="650 439 985 465">Sources: <i>F. martii</i> [13], <i>F. solani</i> [66].</p>
 <p data-bbox="135 677 494 703">(43) Dihydrofusarubin (stereoisomer 1)</p>	<p data-bbox="650 536 935 562">$C_{15}H_{16}O_7$ M_r: 308 mp: 153–154</p> <p data-bbox="650 562 1267 604">UV λ_{max}^{EtOH} nm (log ϵ): 231 (4.17), 243 (4.30), 273 (3.97), 300 (3.71), 391 (3.94).</p> <p data-bbox="650 604 1267 677">1H NMR ($CDCl_3$): 1.75 (<i>s</i>, Me), 3.87 (<i>s</i>, OMe), 6.98 (<i>s</i>, H-Ar), 7.96 (<i>bs</i>, OH), 12.58 (<i>bs</i>, OH), 12.83 (<i>bs</i>, OH), 4.45–4.46 (<i>m</i>, CH), 2.72–1.92 (<i>dd</i>, CH_2), 3.37 (<i>m</i>, 1H), 3.97 (<i>m</i>, 1H).</p> <p data-bbox="650 677 911 703">Source: <i>F. solani</i> [13, 65, 67].</p>
 <p data-bbox="135 915 494 941">(44) Dihydrofusarubin (stereoisomer 2)</p>	<p data-bbox="650 774 935 800">$C_{15}H_{16}O_7$ M_r: 308 mp: 117–118</p> <p data-bbox="650 800 1267 842">UV λ_{max}^{EtOH} nm (log ϵ): 213 (4.15), 243 (4.31), 273 (3.89), 300 (3.70), 391 (3.96).</p> <p data-bbox="650 842 1267 915">1H NMR ($CDCl_3$): 1.38 (<i>s</i>, Me), 3.94 (<i>s</i>, OMe), 6.72 (<i>s</i>, H-Ar), 2.50 (<i>s</i>, OH), 12.34 (<i>s</i>, OH), 2.69 (<i>s</i>, OH), 4.51–4.00 (<i>dd</i>, CH_2), 1.97–1.59 (<i>dd</i>, CH_2), 3.55 (<i>m</i>, 1H), 3.0 (<i>m</i>, 1H).</p> <p data-bbox="650 915 911 941">Source: <i>F. solani</i> [13, 65, 67].</p>
 <p data-bbox="135 1153 424 1179">(45) <i>O</i>-methyldihydrofusarubin</p>	<p data-bbox="650 1012 935 1038">$C_{16}H_{18}O_7$ M_r: 322 mp: 134–135</p> <p data-bbox="650 1038 1267 1080">UV λ_{max}^{EtOH} nm (log ϵ): 242 (4.17), 273 (3.79), 300 (3.7), 390 (3.82).</p> <p data-bbox="650 1080 884 1106">IR ν^{KBr} cm^{-1}: 3425, 1625.</p> <p data-bbox="650 1106 1267 1179">1H NMR ($CDCl_3$): 3.81–2.2 (<i>dd</i>, CH_2), 1.68–2.41 (<i>dd</i>, CH_2), 3.43 (<i>ddd</i>, 1H), 2.94 (<i>ddd</i>, 1H), 1.41 (<i>s</i>, Me), 6.65 (<i>s</i>, H-Ar), 3.95 (<i>s</i>, OMe), 3.23 (<i>s</i>, OMe), 12.02 (<i>s</i>, OH), 12.17 (<i>s</i>, OH).</p> <p data-bbox="650 1179 884 1205">Source: <i>F. solani</i> [13, 65].</p>
 <p data-bbox="135 1399 408 1425">(46) <i>O</i>-ethyl-dihydrofusarubin</p>	<p data-bbox="650 1217 935 1243">$C_{17}H_{20}O_7$ M_r: 336 mp: 138–139</p> <p data-bbox="650 1243 1267 1286">UV λ_{max}^{EtOH} nm (log ϵ): 242 (4.11), 273 (3.74), 300 (3.70), 390 (3.67).</p> <p data-bbox="650 1286 985 1312">IR ν^{KBr} cm^{-1}: 3425 (OH), 1625 (CO).</p> <p data-bbox="650 1312 1267 1384">1H NMR ($CDCl_3$): 3.79–3.4 (<i>dd</i>, CH_2), 1.76–2.45 (<i>dd</i>, CH_2), 3.42 (<i>ddd</i>, 1H), 2.97 (<i>ddd</i>, 1H), 6.65 (<i>s</i>, H-Ar), 1.41 (<i>s</i>, Me), 3.95 (<i>s</i>, OMe), 11.92 (<i>s</i>, OH), 12.16 (<i>s</i>, OH), 5.05 (<i>q</i>, OCH_2CH_3), 1.38 (<i>t</i>, OCH_2CH_3).</p> <p data-bbox="650 1384 1059 1411">Sources: <i>F. solani</i> [13], <i>N. haematococca</i> [68].</p>
 <p data-bbox="135 1636 471 1663">(47) 6-(<i>O</i>)-demethyl-5-deoxyfusarubin</p>	<p data-bbox="650 1475 935 1501">$C_{14}H_{12}O_6$ M_r: 276 mp: 194–196</p> <p data-bbox="650 1501 1267 1544">UV λ_{max}^{MeOH} nm (log ϵ): 2.15 (4.20), 268 (3.89), 288 (3.73), 442 (3.34).</p> <p data-bbox="650 1544 935 1570">IR ν^{KBr} cm^{-1}: 3429, 1649, 1620.</p> <p data-bbox="650 1570 1267 1643">1H NMR ($DMSO-d_6$): 1.43 (<i>s</i>, Me), 6.10 (<i>s</i>, OH), 2.33–2.60 (<i>dd</i>, 1H), 6.93 (<i>d</i>, 1H), 11.20 (<i>s</i>, OH), 6.50 (<i>d</i>, 1H), 12.0 (<i>s</i>, OH), 4.53 (<i>s</i>, 1H).</p> <p data-bbox="650 1643 924 1669">Source: <i>N. haematococca</i> [69].</p>
 <p data-bbox="135 1874 346 1901">(48) Anhydrofusarubin</p>	<p data-bbox="650 1713 935 1739">$C_{15}H_{12}O_6$ M_r: 272 mp: 193–198</p> <p data-bbox="650 1739 911 1766">UV λ_{max}^{EtOH} nm: 235, 290, 537.</p> <p data-bbox="650 1766 1045 1792">IR ν^{KBr} cm^{-1}: 2970, 2920, 2830, 1590, 1435.</p> <p data-bbox="650 1792 1267 1864">1H NMR ($CDCl_3$): 1.95 (<i>s</i>, Me), 3.94 (<i>s</i>, OMe), 5.22 (<i>s</i>, 2H), 5.94 (<i>s</i>, 1H), 6.14 (<i>s</i>, H-Ar), 12.74 (<i>s</i>, OH), 13.10 (<i>s</i>, OH).</p> <p data-bbox="650 1864 1267 1901">Sources: <i>F. solani</i> [7, 56, 65, 70], <i>Neocosmospora vasinfectum</i>, <i>N. africana</i> [57], <i>F. decemcellulare</i> [60].</p>


Table 1—continued.

Naphthoquinone	Same physico-chemical properties
	<p>$C_{15}H_{10}O_7$ M_r: 302 mp: 300° UV λ_{max}^{EtOH} nm: 240, 285, 355, 500. λ_{max}^{MeOH} nm (log ϵ): 255 (4.25), 312 (3.55), 330 (3.46), 508 (3.72), 536 (3.77), 576 (3.72), 612 (3.49). IR ν^{KBr} cm^{-1}: 1745, 1650, 1610, 1582. 1H NMR ($CDCl_3$): 2.40 (s, Me), 3.95 (s, OMe), 6.30 (s, H-Ar), 6.84 (s, 1H), 12.76 (s, OH), 14.30 (s, OH) [71]. Sources: <i>F. solani</i> [71], <i>N. haematococca</i> [72].</p>
(49) Anhydrofusarubin-lacton	
	<p>$C_{15}H_{12}O_7$ M_r: 304 mp: 245° UV λ_{max}^{MeOH} nm (log ϵ): 228 (4.06), 263 (4.05), 278 (4.0), 343 (3.23), 509 (3.80), 535 (3.87), 572 (3.71). IR ν^{KBr} cm^{-1}: 3425, 1600, 1580, 1440. 1H NMR ($CDCl_3$): 2.18 (s, Me), 3.94 (s, OMe), 6.26 (s, 1H), 6.45 (s, 1H), 6.73 (s, 1H), 12.85 (s, OH), 13.25 (s, OH). Sources: <i>F. solani</i> [71], <i>N. haematococca</i> [73].</p>
(50) Anhydrofusarubin lactol	
	<p>$C_{14}H_{10}O_6$ M_r: 274 mp: 202–204 UV λ_{max}^{EtOH} nm: 237, 285, 353, 492sh, 546. IR ν^{KBr} cm^{-1}: 3345, 1660, 1607. 1H NMR ($CDCl_3$) (t): 8.01 (s, Me), 4.79 (s, 2H), 4.03 (s, 1H), 3.69 (s, H-Ar), -1.87 (s, OH), -3.10 (s, OH). Source: <i>Gibberella fujikuroi</i> [74].</p>
(51) O-demethylanhydrofusarubin	
	<p>$C_{15}H_{12}O_5$ M_r: 272 mp: 210–213° UV λ_{max}^{EtOH} nm (log ϵ): 235 (3.39), 295 (5.2), 340 (0.95), 450 (1.02). IR ν^{KBr} cm^{-1}: 1680, 1630. 1H NMR ($CDCl_3$): 1.97 (s, Me), 3.94 (s, OMe), 5.25 (s, 2H), 5.60 (s, 1H), 5.98 (s, H-Ar), 7.12 (s, 1H), 12.2 (s, OH). Source: <i>N. haematococca</i> [62, 70].</p>
(52) Deoxyanhydrofusarubin	
	<p>$C_{14}H_{10}O_5$ M_r: 258 mp: 181–184 UV λ_{max}^{EtOH} nm: 279, 355, 509. IR ν^{KBr} cm^{-1}: 3437, 1644, 1613. 1H NMR ($DMSO-d_6$): 2.03 (s, Me), 5.07 (s, 1H), 5.85 (s, 1H), 6.50 (d, 1H), 6.93 (d, 1H), 11.72 (s, OH), 12.88 (s, OH). Source: <i>N. haematococca</i> [69].</p>
(53) 6-O-demethyl-5-deoxyanhydrofusarubin	
	<p>$C_{16}H_{14}O_6$ M_r: 302 mp: 175–177 UV λ_{max}^{EtOH} nm (log ϵ): 208 (4.3), 222 (4.4), 273 (4.2), 503 (3.6). IR ν^{KBr} cm^{-1}: 3450, 1620, 1570, 1460, 1425. 1H NMR ($CDCl_3$): 2.00 (s, Me), 3.97 (s, OMe), 4.00 (s, OMe), 5.15 (s, 2H), 5.83 (s, 1H), 6.74 (s, 1H), 13.14 (s, OH). Source: <i>F. oxysporum</i> [63].</p>
(54) O-methylanhydrofusarubin	
	<p>$C_{15}H_{14}O_6$ M_r: 290 mp: 207–208 UV λ_{max}^{EtOH} nm: 227, 305, 478sh, 504, 541sh. IR ν^{KBr} cm^{-1}: 1720, 1600. 1H NMR ($CDCl_3$): 2.28 (s, Me), 2.22–3.88 (gh, 2H), 3.91 (s, OMe), 6.18 (s, H), 12.82 (s, OH), 13.21 (s, OH). Sources: <i>F. javanicum</i> [11, 57], <i>F. solani</i> [5, 65], <i>F. martii</i> [5, 12], <i>Neocosmospora vasinfecta</i>, <i>N. africana</i> [57], <i>N. haematococca</i> [56, 70], <i>F. decemcellulare</i> [60].</p>
(55) Javanicin	


Table 1—continued.

Naphthoquinone

Same physico-chemical properties


(56) *O*-methyljavanicin

$C_{16}H_{14}O_6$, M_r : 304 mp: 197–198°
 UV λ_{max}^{MeOH} nm (log ϵ): 226 (4.56), 82.5 (4.04), 482 (3.8), 510 (3.75), 550 (3.80).
 IR ν^{CHCl_3} cm^{-1} : 1620, 1470, 1438, 1270.
 1H NMR ($CDCl_3$): 2.11–3.78 (s, 2H), 2.31 (s, Me), 3.99 (s, OMe), 4.02 (s, OMe), 6.74 (s, H), 13.02 (s, OH).
 Sources: *F. moniliforme* [61], *F. solani* [75], *F. oxysporum* [63].




(57) Deoxyjavanicin

$C_{15}H_{14}O_5$, M_r : 274 mp: 121–124°
 UV λ_{max}^{EtOH} nm (log ϵ): 216 (4.33), 266 (3.97), 290sh (3.70), 428 (3.40).
 IR ν^{KBr} cm^{-1} : 3420, 2940, 2860, 1720, 1665, 1645, 1620, 1580, 1495.
 1H NMR ($CDCl_3$): 2.10 (s, Me), 2.28 (s, Me), 3.77 (s, 2H), 3.87 (s, OMe), 6.60 (s, 1H), 7.20 (s, 1H), 12.28 (s, OH).
 Sources: *Fusarium* sp. [76], *N. haematococca* [70].


(58) *O*-demethyljavanicin

$C_{14}H_{12}O_5$, M_r : 276 mp: 154–155°
 UV λ_{max}^{EtOH} nm: 230, 310, 475, 505, 544.
 IR ν^{KBr} cm^{-1} : 3425, 2940, 2870, 1740, 1635.
 1H NMR ($CDCl_3$): 2.25 (s, Me), 2.29 (s, Me), 3.80 (s, 2H), 3.90 (s, OMe), 6.34 (s, 1H), 13.32 (s, OH), 12.06 (s, OH).
 Source: *F. decemcellulare* [64].




(59) 7-acetyl-5,8-dihydroxy-6-methyl-1,4-NQ

$C_{14}H_{12}O_5$, M_r : 260 mp: 121–122°
 UV λ_{max}^{EtOH} nm: 212, 274, 476, 509, 546.
 IR ν^{KBr} cm^{-1} : 3415, 1715, 1610, 1575, 1465.
 1H NMR ($CDCl_3$): 2.15 (s, Me), 2.32 (s, Me), 3.83 (s, 2H), 7.20 (s, 2H-Ar), 12.42 (s, 1H-OH), 12.57 (s, 1H-OH).
 Source: *Fusarium* sp. [76].




(60) 7-acetyl-5-hydroxy-6-methyl-1,4-NQ

$C_{14}H_{12}O_4$, M_r : 244 mp: 120–122°
 UV λ_{max}^{EtOH} nm: 206, 247, 269, 420, 450.
 IR ν^{KBr} cm^{-1} : 3430, 1720, 1670, 1650, 1625.
 1H NMR ($CDCl_3$): 2.08 (s, Me), 2.27 (s, Me), 3.75 (s, 2H), 7.22 (s, H-Ar), 7.55 (d, 2H-Ar), 12.03 (s, OH).
 Source: *Fusarium* sp. [76].



(61) Anhydrojavanicin

$C_{15}H_{12}O_5$, M_r : 272 mp: 215–217°
 UV λ_{max}^{EtOH} nm: 234, 257, 316, 448.
 IR ν^{KBr} cm^{-1} : 3430, 3130, 1685, 1465.
 1H NMR ($CDCl_3$): 2.37 (s, Me), 2.53 (s, Me), 3.93 (s, OMe), 6.06 (s, 1H), 6.40 (s, H-Ar), 12.85 (s, OH).
 Sources: *Fusarium* sp. [76], *N. haematococca* [56], *F. decemcellulare* [64], *N. vasinfecta* [57], *N. africana* [57].



(62) Dihydroanhydrojavanicin

$C_{15}H_{14}O_5$, M_r : 274 mp: 225–228°
 UV λ_{max}^{EtOH} nm (log ϵ): 225 (4.43), 288 (4.08), 480 (3.81), 501 (3.80), 535 (3.51).
 IR ν^{KBr} cm^{-1} : 3420, 2980, 1670, 1635, 1585.
 1H NMR ($CDCl_3$): 1.40 (d, 3H), 2.13 (s, 3H), 2.50 (dd, 1H), 3.10 (dd, 1H), 3.70 (s, OMe), 5.07 (m, 1H), 6.23 (s, H-Ar), 12.90 (s, OH).
 Sources: *Fusarium* sp. [76], *F. solani* [71].

Table 1—continued.

Naphthoquinone	Same physico-chemical properties
	<p>$C_{15}H_{14}O_6$, M_r: 290 mp: 202–203° UV λ_{max}^{EtOH} nm (log ϵ): 223 (4.06), 297 (3.62), 485 (3.50), 502 (3.47), 538 (3.17). IR ν^{KBr} cm^{-1}: 3490, 1660, 1625, 1585, 1420. 1H NMR ($CDCl_3$): 1.59 (<i>d</i>, Me), 2.67 (<i>t</i>, OH), 2.92 (<i>dd</i>, 1H), 3.46 (<i>dd</i>, 1H), 3.90 (<i>s</i>, OMe), 4.74 (<i>d</i>, 1H), 5.21 (<i>m</i>, 1H), 6.07 (<i>s</i>, H-Ar), 13.60 (<i>s</i>, OH) Source: <i>F. solani</i> [77].</p>
	<p>$C_{14}H_{13}O_6$, M_r: 276 mp: 200–204° UV λ_{max}^{EtOH} nm: 225, 295, 477, 500, 537. IR ν^{KBr} cm^{-1}: 1725, 1610, 1580, 1440, 1360. 1H NMR ($CDCl_3$): 2.28 (<i>g</i>, Me), 3.78 (<i>h</i>, 1H), 7.18 (<i>r</i>, 1H), 3.91 (<i>s</i>, OMe), 6.15 (<i>s</i>, H-Ar), 12.55 (<i>s</i>, OH), 12.59 (<i>s</i>, OH). Sources: <i>Fusarium</i> sp. [78], <i>F. martiela</i> [5, 12], <i>F. solani</i> [65], <i>F. decemcellulare</i> [60], <i>N. vasinfecta</i> [57], <i>N. africana</i> [57], <i>N. haematococca</i> [56].</p>
	<p>$C_{14}H_{12}O_7$, M_r: 292 mp: 231–234° UV λ_{max}^{EtOH} nm (log ϵ): 212 (4.22), 238 (4.14), 261 (4.07), 320 (3.88), 470 (3.84), 490 (3.82), 520 (3.66). IR ν^{KBr} cm^{-1}: 3300, 1710, 1630, 1600, 1570. 1H NMR ($CDCl_3$): 2.30 (<i>s</i>, Me), 3.72 (<i>s</i>, 2H), 4.00 (<i>s</i>, OMe), 6.52 (<i>s</i>, H-Ar), 12.05 (<i>s</i>, OH), 13.20 (<i>s</i>, OH). Sources: <i>F. solani</i> [63], <i>N. haematococca</i> [79].</p>
	<p>$C_{15}H_{14}O_7$, M_r: 306 mp: 226–229° UV λ_{max}^{EtOH} nm (log ϵ): 211 (4.3), 240 (4.2), 270 (4.2), 300 (4.0), 454 (3.9). IR ν^{KBr} cm^{-1}: 3200, 1705, 1635, 1610, 1570. 1H NMR ($CDCl_3$): 2.28 (<i>s</i>, Me), 3.68 (<i>s</i>, 2H), 4.0 (<i>s</i>, OMe), 4.0 (<i>s</i>, OMe), 6.62 (<i>s</i>, H-Ar), 8.21 (<i>s</i>, OH), 13.60 (<i>s</i>, OH). Source: <i>F. oxysporum</i> [63].</p>
	<p>$C_{15}H_{16}O_6$, M_r: 292 mp: 190–194° UV $\lambda_{max}^{dioxane}$ nm (log ϵ): 227 (4.53), 304 (3.97), 500 (3.91). IR ν^{KBr} cm^{-1}: 3350, 3280, 1602. 1H NMR ($CDCl_3$) (τ): 3.85 (<i>s</i>, H-Ar), 6.08 (<i>s</i>, OMe), -3.28 (<i>s</i>, OH), -2.97 (<i>s</i>, OH), 7.67 (<i>s</i>, Me), 7.08 (<i>d</i>, 2H), 5.80 (<i>m</i>, 1H), 6.08 (<i>s</i>, OH), 8.67 (<i>s</i>, Me). Sources: <i>F. solani</i> [80], <i>Fusarium</i> sp. [76].</p>
	<p>$C_{15}H_{16}O_7$, M_r: 308 mp: 206–212° UV λ_{max}^{EtOH} nm (log ϵ): 227 (4.38), 306 (3.87), 452 (3.56), 509 (3.80), 546 (3.59). IR ν^{KBr} cm^{-1}: 3330, 1615, 1550, 1450. 1H NMR ($CDCl_3$): 1.39 (<i>d</i>, Me), 2.86 (<i>dd</i>, 1H), 3.13 (<i>dd</i>, 1H), 3.95 (<i>s</i>, OMe), 4.16 (<i>m</i>, 1H), 4.63 (<i>d</i>, 1H), 4.97 (<i>d</i>, 1H), 6.20 (<i>s</i>, 1H-Ar), 12.86 (<i>s</i>, OH), 13.35 (<i>s</i>, OH). Source: <i>F. solani</i> [77].</p>
	<p>$C_{15}H_{18}O_7$, M_r: 310 mp: 135° UV λ_{max}^{EtOH} nm (log ϵ): 213 (3.98), 244 (4.21), 277 (4.2), 303 (3.64), 391 (3.88), 400 (3.83). IR ν^{KBr} cm^{-1}: 3300, 1610, 1570, 1470, 1430. 1H NMR ($CDCl_3$): 1.22 (<i>d</i>, Me), 1.79 (<i>m</i>, 1H), 1.96 (<i>m</i>, 1H), 2.42 (<i>s</i>, OH), 2.90 (<i>m</i>, 1H), 3.28 (<i>m</i>, 1H), 3.86 (<i>m</i>, 2H), 3.95 (<i>s</i>, Me), 4.24 (<i>dd</i>, 1H), 6.70 (<i>s</i>, H-Ar), 12.22 (<i>s</i>, OH), 12.58 (<i>s</i>, OH). Source: <i>F. solani</i> [77].</p>

Table 1—continued.

Naphthoquinone	Same physico-chemical properties
	<p>$C_{16}H_{18}O_6$ M_r: 306 mp: 152–154° UV λ_{max}^{MeOH} nm (log ϵ): 226 (4.49), 285 (4.02), 476 (3.82), 510sh (3.69). IR ν^{KBr} cm^{-1}: 3460, 1610, 1480, 1435, 1380 [63]. 1H NMR ($CDCl_3$): 1.36 (<i>d</i>, Me), 2.12 (<i>s</i>, Me), 2.62–2.80 (<i>dd</i>, 2H), 3.41 (<i>d</i>, OH), 3.87 (<i>s</i>, OMe), 3.92 (<i>s</i>, OMe), 4.18 (<i>m</i>, 1H), 6.49 (<i>s</i>, H-Ar), 13.22 (<i>s</i>, OH). Sources: <i>Fusarium moniliforme</i> [61], <i>F. solani</i> [63].</p>
	<p>$C_{15}H_{11}NO_5$ M_r: 285 mp: 231–235° UV λ_{max}^{EtOH} nm (log ϵ): 253, 322, 476, 500, 530. IR ν^{CHCl_3} cm^{-1}: 1641, 1591, 1311. 1H NMR ($CDCl_3$): 2.81 (<i>s</i>, Me), 7.96 (<i>s</i>, 1H), 4.02 (<i>s</i>, OMe), 9.50 (<i>s</i>, 1H), 6.75 (<i>s</i>, H-Ar), 13.10 (<i>s</i>, OH), 13.50 (<i>s</i>, OH). Sources: <i>F. bostrycoides</i> [81], <i>F. solani</i> [65, 82], <i>F. decemcellulare</i> [60].</p>
	<p>$C_{16}H_{13}NO_5$ M_r: 299 mp: 215–216° UV λ_{max}^{MeOH} nm (log ϵ): 247 (4.50), 318 (3.92), 480 (3.83). λ_{max}^{Me-HCl} nm (log ϵ): 227 (4.32), 262 (4.20), 310 (3.88), 510 (3.73). λ_{max}^{Me-ONa} nm (log ϵ): 259 (4.44), 306 (3.82), 546 (4.0). IR ν^{KBr} cm^{-1}: 1641, 1591, 1311. 1H NMR ($CDCl_3$): 9.44 (<i>s</i>, 1H), 7.85 (<i>s</i>, 1H), 6.86 (<i>s</i>, H-Ar), 2.76 (<i>s</i>, Me), 4.05 (<i>s</i>, OMe), 4.05 (<i>s</i>, OMe). Sources: <i>F. moniliforme</i> [61], <i>F. oxysporum</i> [63].</p>
	<p>$C_{14}H_9NO_5$ M_r: 271 mp: 221–223° UV λ_{max}^{EtOH} nm: 252, 340, 476, 496, 530. λ_{max}^{Et-ONa} nm: 230, 290, 485sh, 525, 556. Source: <i>F. decemcellulare</i> [64].</p>
	<p>$C_{17}H_{11}NO_4$ M_r: 269 mp: 195–196° UV λ_{max}^{MeOH} nm: 237, 270, 322(sh), 414. IR ν^{KBr} cm^{-1}: 1680, 1635, 1590. 1H NMR ($CDCl_3$): 2.78 (<i>s</i>, Me), 3.94 (<i>s</i>, OMe), 6.74 (<i>d</i>, 1H), 7.32 (<i>d</i>, 1H), 7.86 (<i>s</i>, 1H), 9.40 (<i>s</i>, 1H), 12.76 (<i>s</i>, OH). Source: <i>N. haematococca</i> [83].</p>
	<p>$C_{16}H_7NO_4$ M_r: 255 mp: 300–305° UV λ_{max}^{MeOH} nm (log ϵ): 205 (4.40), 239 (4.42), 280 (4.12), 320 (sh, 3.83), 421 (3.75). IR ν^{KBr} cm^{-1}: 3200–2400, 1682, 1637, 1600. 1H NMR [(CD_3)$_2$SO]: 2.70 (<i>s</i>, Me), 6.60 (<i>d</i>, 1H), 7.10 (<i>d</i>, 1H), 7.80 (<i>s</i>, 1H), 9.20 (<i>s</i>, 1H), 12.57 (<i>s</i>, OH). Source: <i>N. haematococca</i> [84].</p>
	<p>$C_{15}H_{12}O_6$ M_r: 302 mp: 200–210° IR ν^{KBr} cm^{-1}: 3300–2500. 1H NMR (CD_3OD): 2.25 (<i>s</i>, Me), 3.98 (<i>s</i>, 2H), 4.04 (<i>s</i>, OMe), 6.46 (<i>s</i>, H-Ar). Source: <i>N. haematococca</i> [85].</p>

Table 1—continued.









Naphthoquinone	Same physico-chemical properties
	<p>$C_{15}H_{14}O_7$, M_r: 306 mp: 162° UV λ_{max}^{EtOH} nm: 302, 477sh, 505, 542. Sources: <i>F. solani</i>, <i>F. martii</i> [12], <i>Neocosmospora vasinfectum</i>, <i>N. africana</i> [57], <i>F. decemcellulare</i> [60].</p>
(77) Novarubin	
	<p>$C_{18}H_{16}O_9$, M_r: 376 mp: 180–182 UV λ_{max}^{EtOH} nm: 229, 304, 474, 501, 536. IR ν^{KBr} cm^{-1}: 3600–2800, 1720. 1H NMR ($CDCl_3$): 1.68 (s, Me), 2.73 (d, 1H), 3.12 (d, 1H), 2.10 (m, 1H), 5.42 (d, 1H), 2.86 (m, OH), 3.94 (s, OMe), 4.58 (m, 1H), 12.53 (s, OH), 12.93 (s, OH). Sources: <i>F. solani</i> [65], <i>F. martii</i> [5, 12].</p>
(78) Marticin	
	<p>$C_{18}H_{16}O_9$, M_r: 376 mp: 160–163 UV λ_{max}^{EtOH} nm: 229, 304, 474, 501, 536. IR ν^{KBr} cm^{-1}: 3430, 1720. 1H NMR ($CDCl_3$): 1.70 (s, Me), 3.00 (d, 1H), 3.13 (d, 1H), 2.04 (m, 1H), 5.58 (d, 1H), 2.30 (m, OH), 4.31 (dd, 1H), 6.20 (s, H-Ar), 12.51 (s, OH), 12.91 (s, OH). Sources: <i>F. solani</i> [65], <i>F. martii</i> [5, 12].</p>
(79) Isomarticin	
	<p>$C_{15}H_{12}O_7$, M_r: 304 mp: 230 UV λ_{max}^{EtOH} nm: 255, 320, 443, 465. IR ν^{KBr} cm^{-1}: 3350, 3250–2500, 2840, 1735, 1605. 1H NMR ($CDCl_3$): 1.67 (s, Me), 3.92 (s, OMe), 5.18 (q, 1H), 6.66 (s, H-Ar), 8.10 (s, 1H), 13.03 (s, OH), 13.35 (s, OH). Sources: <i>N. haematococca</i> [72], <i>F. oxysporum</i> [14].</p>
(80) Nectriafurone	
	<p>$C_{16}H_{14}O_7$, M_r: 318 mp: 214–222 UV λ_{max}^{EtOH} nm: 236 (4.37), 225sh (4.21), 322 (3.90), 443 (4.02). IR ν^{KBr} cm^{-1}: 3430, 3120, 1645, 1625, 1600. 1H NMR ($CDCl_3$): 1.64 (d, Me), 4.00 (s, OMe), 4.03 (s, OMe), 4.78 (d, OH), 5.19 (m, 1H), 6.83 (s, H-Ar), 7.99 (s, 1H), 13.30 (s, OH). Source: <i>F. oxysporum</i> [14].</p>
(81) O-methyl-nectriafurone	
	<p>$C_{20}H_{14}O_8$, M_r: 382 mp: 320–325 (decomp.) UV $\lambda_{max}^{CHCl_3}$ nm (log ϵ): 253 (4.53), 271 (4.51), 320 (3.92), 510 (3.96), 550 (3.77). IR ν^{KBr} cm^{-1}: 1670, 1653, 1620. 1H NMR ($CDCl_3$): 3.07 (3H, s-Me), 4.23 (s, 2OMe), 6.93 (1H, s-Ar), 7.44 (1H, m), 7.49 (1H, d). Sources: <i>Gibberella fujikuroi</i> [17, 89], <i>Fusarium bulbigenum</i> [86], <i>Fusarium oxysporum</i> [87], <i>Verticillium agaricinum</i> [88].</p>
(82) Bikaverin	
	<p>$C_{19}H_{12}O_8$, M_r: 368 mp: > 300° (decomp.) UV $\lambda_{max}^{CHCl_3}$ nm (log ϵ): 253 (4.51), 273 (4.51), 320 (3.92), 515 (3.96), 550 (3.78). IR ν^{Nujol} cm^{-1}: 3400, 1670, 1640, 1620–1560. 1H NMR ($CDCl_3$): 3.02 (s, Me), 4.18 (s, OMe), 6.92 (s, 1H), 7.26 (m, 1H), 7.45 (d, 1H). Source: <i>Gibberella fujikuroi</i> [89].</p>
(83) Norbikaverin	

Table 1—continued.


Naphthoquinone

Same physico-chemical properties




(84) Biflaviolin

$C_{25}H_{20}O_{10}$ M_r : 410
 UV λ_{max}^{EtONa} nm (log ϵ): 294 (4.68), 372 (4.02), 422sh, 560 (3.43).
 λ_{max}^{Et-HCl} nm (log ϵ): 265 (4.48), 312 (4.15), 390 (3.81), 460sh.
 1H NMR [(CD₃)₂CO]: 6.68 (d, 2H), 7.18 (d, 2H), 12.5 (s, 2OH).
 Source: *Thielaviopsis basicola* [90].




(85) Xanthomegnin

$C_{30}H_{26}O_{12}$ M_r : 578 mp: > 260° (dec.)
 UV λ_{max}^{MeONa} nm (log ϵ): 222 (4.41), 264 (4.29), 380 (3.90).
 IR ν^{KBr} cm⁻¹: 3430, 1721, 1675, 1616, 1590.
 1H NMR (CDCl₃): 1.51 (d, Me), 3.02 (d, CH₂), 4.67 (m, 1H), 7.50 (s, 1H), 4.11 (s, OMe), 13.17 (s, OH).
 Sources: *Trichophyton rubrum* [91], *T. megnini* [92], *T. violaceum* [93], *Aspergillus sulphureus*, *A. melleus* [94], *Penicillium viridicatum* [95], *Microsporium cookei* [96], *Nannizzia cajetani* [97].




(86) 3,4-dehydroxanthomegnin

$C_{30}H_{24}O_{12}$ M_r : 576 mp: > 160° (dec.)
 UV λ_{max}^{MeOH} nm: 237, 275sh, 385, 436sh.
 λ_{max}^{MeONa} nm: 254, 275, 300, 394, 550.
 IR ν^{KBr} cm⁻¹: 3385, 1738, 1680, 1655, 1618.
 1H NMR (CDCl₃) (quinone part): 1.56 (d, Me), 4.68 (m, 1H), 3.05 (d, 2H), 7.51 (s, 1H), 4.16 (s, OMe), 13.17 (s, H); (phenolic part): 2.35 (s, Me), 6.39 (s, 2H), 7.48 (s, 1H), 4.16 (s, OMe), 13.7 (s, Me).
 Sources: *Nannizzia cajetani* [97], *P. citreoviride* [98].




(87) 3,4,3',4'-Bisdehydroxanthomegnin

$C_{30}H_{22}O_{12}$ M_r : 574
 UV λ_{max}^{MeOH} nm: 231sh, 267sh, 364.
 λ_{max}^{MeONa} nm: 229, 282, 387, 530.
 1H NMR (CDCl₃): 2.37 (s, Me), 6.39 (s, 2H), 7.53 (s, 1H), 4.19 (s, OMe), 13.72 (s, OH).
 Source: *N. cajetani* [97].



(88) Viomellein

$C_{30}H_{26}O_{11}$ M_r : 562 mp: > 260° (dec.)
 UV λ_{max}^{MeOH} nm: 264, 366sh, 376.
 λ_{max}^{MeONa} nm: 265, 330, 372, 385, 536.
 IR ν^{KBr} cm⁻¹: 3445, 1735, 1675, 1640.
 1H NMR (CDCl₃) (quinone part): 1.54 (s, Me), 4.70 (m, 1H), 3.01 (d, 2H), 7.49 (s, 1H), 3.90 (s, OMe), 13.50 (s, OH); (phenolic part): 1.54 (s, Me), 4.70 (m, 1H), 3.01 (d, 2H), 6.96 (s, 1H), 6.66 (s, 1H), 3.86 (s, OMe), 9.80 (OH), 13.88 (OH).
 Sources: *N. cajetani* [97], *A. sulphureus*, *A. melleus* [94], *T. violaceum* [93], *P. viridicatum* [95], *P. citreoviride* [98].



(89) 3',4'-dehydroviomellein

$C_{30}H_{24}O_{11}$ M_r : 560 mp: > 150° (dec.)
 UV λ_{max}^{MeOH} nm: 261, 365, 375, 446.
 λ_{max}^{MeONa} nm: 260, 372, 383, 547.
 IR ν^{KBr} cm⁻¹: 3390, 1752, 1675, 1648, 1618.
 1H NMR (CDCl₃) (quinone part): 2.33 (s, Me), 6.33 (s, 2H), 7.53 (s, 1H), 3.92 (s, OMe), 14.03 (s, OH); (phenolic part): 1.56 (d, Me), 4.72 (m, 1H), 3.02 (d, 2H), 6.96 (s, 1H), 6.98 (s, 1H), 3.86 (s, OMe), 9.80 (s, OH), 13.88 (s, OH).
 Source: *N. cajetani* [97].


Table 1—continued.

Naphthoquinone	Same physico-chemical properties
	<p>$C_{26}H_{20}O_9$ M_r: 476 mp: 133–136° UV λ_{max}^{MeOH} nm (log ϵ): 225 (4.06), 264 (4.3), 384 (3.91). IR ν^{KBr} cm^{-1}: 3420, 1725, 1590. 1H NMR ($CDCl_3$): 1.50 (<i>d</i>, Me), 2.98 (<i>d</i>, 2H), 3.83 (<i>s</i>, OMe), 3.89 (<i>s</i>, OMe), 4.74 (<i>m</i>, 1H), 6.66 (<i>s</i>, 1H-Ar), 6.95 (<i>s</i>, 1H-Ar), 9.78 (<i>s</i>, OH), 12.30 (<i>s</i>, OH), 13.92 (<i>s</i>, OH). Source: <i>Penicillium veridicatum</i> [99].</p>
(90) Xanthoviridicatin D	
	<p>$C_{25}H_{16}O_8$ M_r: 444 mp: 318–320° UV λ_{max}^{MeOH} nm (log ϵ): 280 (4.47), 355 (3.51), 440 (3.64). IR ν^{KBr} cm^{-1}: 3390, 1737, 1664, 1580. 1H NMR ($CDCl_3$): 1.53 (<i>d</i>, Me), 2.98 (<i>d</i>, 2H), 4.10 (<i>s</i>, OMe), 4.72 (<i>m</i>, 1H), 6.87 (<i>s</i>, H-Ar), 7.37 (<i>s</i>, H-Ar), 7.44 (<i>dd</i>, H-Ar), 7.80 (<i>t</i>, H-Ar), 8.19 (<i>dd</i>, H-Ar), 12.30–13.92 (<i>s</i>, 2OH). Source: <i>Penicillium veridicatum</i> [99].</p>
(91) Xanthoviridicatin G	
	<p>$C_{25}H_{20}O_{10}$ M_r: 528 mp: 280 UV $\lambda_{max}^{CHCl_3}$ nm (log ϵ): 280 (4.50), 357 (3.58), 415 (3.69). IR ν^{KBr} cm^{-1}: 3278, 2910, 1729, 1670, 1640, 1600. Sources: <i>P. veridicatum</i> [95], <i>A. sulphureus</i>, <i>A. nulleus</i> [94].</p>
(92) Rubrosulphin	
	<p>$C_{28}H_{18}O_{12}$ M_r: 546 Source: <i>Microsporium cookei</i> [100].</p>
(93) Luteosporin	
	<p>$C_{30}H_{18}O_{12}$ M_r: 570 mp: >330° (dec.) UV $\lambda_{max}^{CHCl_3}$ nm (log ϵ): 248 (4.62), 268 (4.48), 381 (3.94). IR ν^{KBr} cm^{-1}: 1660, 1600, 1470, 1438. 1H NMR (CF_3COOD) (τ): 7.13 (<i>s</i>, Me), 5.63 (<i>s</i>, OMe), 1.87 (<i>s</i>, H-Ar), 2.91 (<i>s</i>, 2H). Sources: <i>Hypomyces rosellus</i> [101], <i>Dactylium dendroides</i> [101], <i>F. culmorum</i> [102, 103], <i>F. graminearum</i> [102, 104, 105], <i>F. decem-cellulare</i> [106, 107].</p>
(94) Aurofusarin	
	<p>$C_{30}H_{20}O_{10}$ M_r: 556 mp: >300° UV λ_{max}^{EtOH} nm (log ϵ): 225 (4.50), 281 (4.63), 346 (3.89), 405 (4.01). IR ν^{CHCl_3} cm^{-1}: 1665, 1650, 1620, 1603. 1H NMR ($CDCl_3$) (τ): 7.76 (<i>s</i>, MeCOO), 7.60 (<i>s</i>, MeCOO, Me, Me), 6.10 (<i>s</i>, OMe), 3.97 (<i>s</i>, H), 3.86 (<i>s</i>, H), 3.05 (<i>s</i>, H), 2.93 (<i>s</i>, H), 1.90 (<i>s</i>, H), 4.74 (<i>s</i>, OH). Source: <i>Fusarium culmorum</i> [106].</p>
(95) Fuscofusarin	


Table I—continued.

Naphthoquinone


Same physico-chemical properties




(96) Purpurogenone

 $C_{29}H_{20}O_{11}$ M_r : 544 mp: 310° (dec.)UV $\lambda_{max}^{CHCl_3}$, nm (log ϵ): 252 (4.18), 306 (3.73), 387 (3.80), 498 (3.42), 528 (3.47), 568 (3.27).Source: *Penicillium purpurogenum* [108].

(97) Viopurpurin

 $C_{29}H_{20}O_{11}$ M_r : 544 mp: 310°UV $\lambda_{max}^{CHCl_3}$, nm (log ϵ): 272 (4.58), 280 (4.59), 375 (3.92).IR ν^{KBr} cm^{-1} : 3420, 1730, 1670, 1635, 1600, 1560, 1545.Sources: *P. viridicatum* [95], *T. violaceum* [93], *A. melleus*, *A. sulphureus* [94].


(98) Xylindein

 $C_{32}H_{24}O_{10}$ M_r : 564 mp: > 300°UV λ_{max}^{EtOH} nm: 380, 405, 423, 603, 647.IR ν^{KBr} cm^{-1} : 1720, 1625.Sources: *Chlorociboria aeruginosa*, *Lophyostoma viridarium* [109].

(99) Mycochrison

 $C_{20}H_{12}O_8$ M_r : 364 mp: 195° (dec.)UV λ_{max}^{EtOH} nm (log ϵ): 236 (4.49), 303 (4.75), 438 (3.80).IR ν^{KBr} cm^{-1} : 1689, 1666, 1637. 1H NMR [(CD_3) $_2CO$] (τ): 5.88 (*m*, 1H), 6.38 (*m*, 1H), 3.5 (*d*, 1H), 3.6 (*s*, 1H), 3.57 (*d*, 1H), 3.78 (*d*, 1H).

Source: cultures an unidentified inopercolate discomycete [110].



(100) Dermocanarin 2

 $C_{33}H_{26}O_{11}$ M_r : 598 mp: 235–240°UV λ_{max}^{EtOH} nm (log ϵ): 216 (4.80), 272 (4.51), 302sh (4.21), 420 (3.82). λ_{max}^{EtOH} nm (log ϵ): 515 (3.75).IR ν^{KBr} cm^{-1} : 3457, 1771, 1651, 1633, 1593. 1H NMR ($CDCl_3$): 7.12 (*br s*, 1H), 7.64 (*br s*, 1H), 7.83 (*s*, 1H), 2.47 (*s*, Me), 3.98 (*s*, OMe), 12.38 (*s*, OH), 2.47 (*d*, 1H^A), 1.97 (*d*, 1H^B), 2.71 (*d*, 1H^B), 3.36 (*d*, 1H^B), 7.32 (*d*, 1H), 6.75 (*d*, 1H), 1.39 (*s*, Me), 3.03 (*s*, OH), 3.96 (*s*, OMe), 3.92 (*s*, OMe).Source: *Dermocybe canaria* [111].

UV-wavelength are in nm. and log ϵ or ϵ values are given in parentheses. Only values for λ_{max} are quoted. IR values are in cm^{-1} . Melting points (MP) are in degrees centigrade and are uncorrected. 1H NMR chemical shifts are in ppm on the δ scale.

Table 2. Fungi producers of naphthoquinone metabolites

Culture of fungi	Metabolites (N in Tab. 1)	Culture of fungi	Metabolites (N in Tab. 1)
1. <i>Aspergillus citricus</i>	5	33. <i>Hypomyces rosellus</i>	94
2. <i>A. melleus</i>	85, 88, 92, 97	34. <i>Lambertella</i> spp.	25
3. <i>A. niger</i>	5	35. <i>Lophyostoma viridarum</i>	98
4. <i>A. parvulus</i>	6	36. <i>Marasmius graminum</i>	1
5. <i>A. sulfureus</i>	85, 88, 92, 97	37. <i>Microsporium cookei</i>	85, 93
6. <i>Cercospora melonis</i>	7	38. <i>Metatrichia vespasiarum</i>	17, 18
7. <i>Cetraria cucullata</i>	20	39. <i>Mollisia caesia</i>	14, 15
8. <i>Chlorociboria aeruginosa</i>	98	40. <i>Mollisia fallens</i>	15
9. <i>Cladosporium</i> sp.	3, 4, 5	41. <i>Nannizzia cajetani</i>	81, 85-89
10. <i>Cladonia</i> spp.	28	42. <i>Neocosmospora africana</i>	37, 48, 55, 61, 64, 77
11. <i>Corinespora custiocola</i>	9, 10	43. <i>Neocosmospora vasinfectum</i>	37, 48, 55, 61, 64, 77
12. <i>Criptosporium pinicola</i>	27	44. <i>Nectria haematococca</i>	37, 38, 46, 47, 49, 50, 52, 53, 55, 57, 61, 64, 65, 74-76, 80
13. <i>Cylindrocarpon</i> sp.	19	45. <i>Penicillium</i> spp.	16
14. <i>Dactylium dendroides</i>	94	46. <i>P. ceterioviride</i>	86, 88
15. <i>Dermocybe canaria</i>	100	47. <i>P. purpurogenum</i>	96
16. <i>Foma vasabiae</i>	8	48. <i>P. viridicatum</i>	85, 88, 80, 97
17. <i>Fomes annosus</i>	30	49. <i>Pirenochetia cerestris</i>	31, 32
18. <i>Fusarium bostricoideas</i>	71	50. <i>Pirex coccifera</i>	28
19. <i>F. bulbigenum</i>	82	51. <i>Pseudospirones simplex</i>	25
20. <i>F. culmorum</i>	94, 95	52. <i>Stemphyllium</i> spp.	3, 4, 5
21. <i>F. decemcellulare</i>	37, 40, 48, 58, 61, 62, 73, 77, 94	53. <i>Thielaviopsis basicola</i>	84
22. <i>F. graminearum</i>	94	54. <i>Torula herbarum</i>	22, 23, 24
23. <i>F. javanicum</i>	55	55. <i>Trichia floriformis</i>	17, 18
24. <i>F. moniliforme</i>	37, 39, 56, 70	56. <i>Trichophyton megnini</i>	85
25. <i>F. martii</i>	12, 13, 64	57. <i>T. rubrum</i>	85
26. <i>F. oxisporum</i>	39, 54, 56, 66, 80, 82	58. <i>T. violaceum</i>	85, 88, 97
27. <i>F. solani</i>	37, 42-50, 55-71, 79-79	59. <i>Ushnea canariensis</i>	29
28. <i>Gibberella fujicuroi</i>	51, 82, 83	60. <i>Ushnea nookeri</i>	29
29. <i>Gnomonia eristoma</i>	33, 34, 35, 36	61. <i>Ustilago</i> spp.	26
30. <i>Helicobasidium mompa</i>	13	62. <i>Verticillium agaricinum</i>	82, 83
31. <i>Hematoma ventosum</i>	31	63. <i>Verticillium dahliae</i>	2, 3, 4, 5
32. <i>Hendersonula toruloidea</i>	11, 12		

nitrogen led to the inhibition of naphthoquinone biosynthesis.

Kurobane *et al.* [82] noted that the composition of naphthoquinones can be determined by the initial ratio of the carbon and nitrogen sources in the culture medium. Thus, when grown on maltose (at a concentration of 20–50 g/l) the fungus *F. solani* synthesized dihydrofusarubins and javanicin if ammonium tartrate was added at 4.6 g/l. An increase in the concentration of the nitrogen source to 6.9 g/l led to the synthesis of bostricoidin, the molecule of which contains a nitrogen atom. A significant role in the increase of pigment formation was played by Fe, Mg and Zn ions [112]. A study of the effect of the cultivation conditions on the biosynthesis of naphthoquinones by the fungus *F. decemcellulare*, showed that the major condition of pigment formation is the inhibition or total cessation of the fungal growth with an excess carbon source and energy [107]. The factor regulating the composition of naphthoquinones proves to be the pH of the cultivation medium. Thus,

during the inhibition of fungal growth by a high concentration of hydrogen ions in the medium (pH 4.0 and lower) and excess carbon, we observed synthesis of naphthasarins (fusarubin, javanicin, bostricoidin etc.) [107].

On the other hand, inhibition of fungal growth as a result of a pH increase in the medium to 8.0 was accompanied by the formation of only the dimeric naphthoquinone, aurofusarin [107]. Formation of aurofusarin was also observed during the limitation of fungal growth by nitrogen and phosphorous sources (optimal pH being maintained). It proved that during the growth on a medium containing ammonium sulphate as a nitrogen source there occurred the physiological acidification of the cultivation medium to pH 4.0 and lower, due to the consumption of ammonium and accumulation of the acid. Under these conditions, naphthasarins metabolites were formed.

In contrast, in growth with sodium nitrate (Chapek's medium), the pH of the medium was increased to 8.0 and aurofusarin accumulated. In the

limitation of fungal growth by a phosphorous source at excess ammonium sulphate and the optimal pH, dimeric naphthoquinone, aurofusarin was formed and no synthesis of naphthasarins was observed. Based on these data, it was concluded that pH of the medium, but not the nature of the nitrogen source, determined the composition of naphthoquinones.

A pH decrease as a necessary factor of naphthoquinone synthesis was earlier reported by Backer *et al.* [6], Claydon *et al.* [9] and Kurobane *et al.* [82]. The key role of pH in naphthoquinone synthesis is probably due to the fact that at pHs close to the neutral value, naphthoquinones exhibit a strong cytotoxic action against the microorganisms, plants and the fungal producers themselves [60, 113–115].

In contrast, at low pH the activity of the pigments, as auto-oxidative compounds, sharply decreases [60]. We suggest that, depending on the ambient conditions, the fungal producers are capable of changing the nature of the end product synthesized, thus protecting themselves from the harmful effects of their own metabolites.

Among other factors which lead to the synthesis of naphthoquinones, we should note the following [56]: (1) the presence of inhibitors capable of inhibiting the growth of the fungi (5-fluorouracil, ethidium bromide, sodium azide, nystatin); (2) temperature not optimal for the growth (30–32° instead of 26°); (3) oxidation of the incubation medium (pH lower than 3.5); and (4) the presence of microbial antagonists (*Bacillus subtilis*) or their products.

All the above enables one to consider naphthoquinones of the fungi (exemplified by the genus *Fusarium*) as a classical example of the secondary metabolites which are synthesized under conditions of growth inhibition or total cessation of growth [116].

THE MECHANISM OF NAPHTHOQUINONE SYNTHESIS

Detailed studies of the mechanism of naphthoquinone synthesis by fungi began in 1965 [117, 118]. The authors using ¹⁴C-labelled acetate showed the synthesis of javanicin by the fungus *F. javanicum* to proceed via the polyketide route. Experiments with the ¹⁴C-labelled methyl group of methionine showed that the methoxy group of javanicin originated from this source as a result of the interaction of the alkylating agent *S*-adenosylmethionine with the aromatic nucleus. Formation of the methyl group of the pigment occurs as a result of the reduction of the end carboxyl group. Subsequently, these results were supported by the data of NMR spectroscopy obtained by Kurobane *et al.* [37] in studies of dihydrofusarubin synthesis by *F. solani* using ¹⁴C- and ³H-labelled acetate compounds. Similar data were obtained in studies of 5-deoxyfusarubin synthesis by a mutant strain *N. haematococca* by Parisot *et al.* [62] and by Holenstein [117] for marticin synthesis. The data cited indicate that synthesis of naphthoquinones by fungi proceeds via the formation of a common precursor—a product

of the acetate malonate pathway. Figure 1 summarizes the data on the biogenic relations of various naphthoquinones synthesized by *Fusarium* fungi [37, 80, 82, 85, 118].

Gatenbeck and Bentley [118] and later Arsenault [80] suggested that the primary metabolite in pigment synthesis is an aromatic acid (Fig. 1), which is then methylated to fusarubinic acid. This metabolite was not found for over 20 years. Only in 1988, was it isolated from the culture medium of *N. haematococca* and characterized [85]. As a result of successive reduction, fusarubinic acid was subsequently transformed to the aromatic aldehyde (not yet found), to the primary alcohol fusarubin and then to javanicin, solaniol and bostricoidin [80, 118]. Anhydrofusarubin and anhydrojavanicin were formed as a result of dehydration of, respectively, fusarubin and javanicin. Fusarubinic acid can also be directly converted into anhydrofusarubin lactone which, in turn, is reduced to anhydrofusarubin lactol [3, 85].

A slightly different scheme of naphthoquinone biogenesis was proposed by Kurobane *et al.* [37, 82]. They postulated that the next product synthesized after the aromatic acid is dihydrofusarubin (4). Under alkaline conditions, dihydrofusarubin is converted to fusarubin as a result of nonenzymatic oxidation. Similarly, norjavanicin and bostricoidin are formed (for bostricoidin to be formed, ammonium ions have to be present in the medium). Javanicin found in the medium at the early stages of cultivation is believed by the authors [82] to be a co-metabolite of dihydrofusarubin, but not a product of its conversion.

The polyketide route of synthesis has been also demonstrated for other metabolites: mollisin [36], mompain [59], fomasarin [52], flaviolin [27].

THE BIOLOGICAL ACTIVITY OF NAPHTHOQUINONE METABOLITES

As we noted in the Introduction, naphthoquinones have a broad range of biological action. The fungal pigments are found to be active against bacteria, yeasts, fungi [5, 10–13, 15, 119–121], protozoa *Leishmania brasiliensis* [17], insects *Calliphora erythrocephala* [9]. The cytotoxic activity of naphthoquinones against mouse leukemia [13] and HeLa cells [15] has been noted. Along with the antibiotic and toxic activities, naphthoquinones revealed mutagenic and carcinogenic properties [122].

One should note that the antibiotic activity of naphthoquinones is of selective character and is manifest mainly against Gram-positive bacteria. This effect is suggested [15] to be due to the lipophilic properties of naphthoquinones and their inability to penetrate the outer membrane of Gram-positive bacteria. This is supported by a relatively strong inhibitory effect of naphthoquinones against *Escherichia coli*, the permeability barrier being decreased for antibiotics [15].

Of special interest from the biological and medical points of view are dimeric naphthoquinones of the




Fig. 1. Proposed route for the formation of naphthoquinone metabolites in fungi genus *Fusarium*. 1—aromatic acid; 2—fusarubicin acid; 3—aromatic aldehyde; 4—dihydrofusarubin; 5—fusarubin; 6—javanicin; 7—solaniol; 8—anhydrofusarubin; 9—norjavanicin; 10—bostricoidin; 11—anhydrojavanicin; 12—anhydrofusarubin lacton; 13—anhydrofusarubin lactol.

fungi. Besides the antibiotic activity against Gram-positive and Gram-negative bacteria, fungi and yeasts [97, 98, 107], many of them exhibit the properties of mycotoxins [123]. Thus, the best known xanthomegnin and viomellein, synthesized by *Penicillium*, *Aspergillus*, *Microsporium*, *Trichophyton* fungi, affected liver and kidney in laboratory animals [123].

Most naphthoquinone metabolites are synthesized

by phytopathogenic fungi *Fusarium*, and their phytotoxic effects have been studied the most intensively [5–8, 12, 57, 75, 76, 124, 128].

Already the first studies on pea seedlings showed the symptoms of a disease during the treatment of the seedlings with solutions of purified naphthoquinones [5, 8, 12]. Subsequently, naphthoquinones were found to be able to inhibit the growth of the roots of the

radish, lemon [6, 8], and lettuce seedlings [75], to suppress the growth of the tomato culture *Lycopersicon esculentum* [39], to slow down the germination of the pine-tree (*Pinus thunbergii*) pollen [76].

Treatment of the developing rough lemon seedlings with naphthoquinones caused leaf roll, which occurred between 48–72 hr, followed by leaf dehydration, wilt and veinal chlorosis [8]. Characteristically, the pea and lemon sprouts infected by *F. solani* isolates, exhibited similar features [8]. Interestingly, in the incubation of the pea and lemon seedlings in a medium with isomarticin and fusarubin, these pigments were found in stems and leaves in 4 hr [8, 124].

The phytotoxic activity as the antibiotic effects, depended on the structure of the naphthoquinones. Marticin and isomarticin were more toxic to peas and lemon than fusarubin, javanicin, norjavanicin and anhydrofusarubin [5, 8].

THE MECHANISM OF TOXIC ACTION

Studies of the mechanism of the antibiotic and phytotoxic action of naphthoquinones were started by Kern *et al.* [5, 125, 126]. Experiments on the action of the pigments on the subcellular level showed [125] that the naphthoquinone metabolites inhibited the aerobic decarboxylation of α -ketoglutarate and the anaerobic decarboxylation of pyruvate. Since the inhibitory effect was partially eliminated in the presence of excess thiamine pyrophosphate (TPP), the authors concluded that the pigments reacted directly with TPP.

The toxicity of the pigments for bacteria and plants was found to be decreased in the presence of some metal ions (Cu, Fe, Al) [5, 126, 127]. Thus, the presence of fusarubin and javanicin in the medium with the increased amount of Cu had no effect on the growth of plants and bacteria. The increase in the content of Cu ions in the growth medium did not affect the formation of the pigments [5, 127]. This effect of detoxication of naphthoquinones, in the opinion of the authors, was a result of formation of chelate complexes with metal ions. It is not impossible that the ability to form complexes with metals also determines their toxicity.

Further studies have found that the toxic effect of marticins on the development of pea sprouts decreases in the presence of glutamine [128]. It was suggested that the toxins inhibit glutamine synthetase. This hypothesis was supported in *in vitro* experiments on the direct inhibition of glutamine synthetase as well as on the reconstitution of the effect of isomarticin by nethionine sulphoximine, a known inhibitor of glutamine synthetase [5, 128].

It was shown [5, 13, 125] that the antibacterial activity of the fungal pigments depended on their chemical structure. Thus, in the metabolites synthesized by *Fusarium solani*, the activity against *Bacillus subtilis* decreased as follows: novarubin > norjavanicin > javanicin > fusarubin > marticin [5].

The activity of novarubin exceeded that of marticin ten-fold. The fungicidal activity changed in the same order but differed a 100-fold [5].

The dependence of the activity on the structure of a metabolite was also shown by Kurobane *et al.* [13]. Dihydrofusarubin proved to be more active than fusarubin; in turn, dihydrofusarubin and fusarubin were more active than their *O*-methyl and ethyl derivatives. The authors believe this to be due to the decrease in their solubility in water [13].

Besides the above effects, marticin and isomarticin evoked a pronounced disturbance of the selective permeability of plant cell membranes. Similar changes were observed in pea cells infected by a pathogenic strain *F. martii* which synthesizes large amounts of marticin and isomarticin. In both cases, the cytoplasmic membrane and chloroplast membrane were disrupted which led to the release of the nucleotides, amino acids, chlorophyll, proteins and mineral salts from the cells [5, 57, 128].

This picture of the consequences of the effect of naphthoquinones on the plant and microbial cells may reflect their ability to uncouple oxidative phosphorylation. It is just the property of uncoupling oxidative phosphorylation that is, in the opinion of the Japanese investigators, a cause of the high antibiotic activity of simple and dimeric naphthoquinones [100, 129, 130]. It was shown that a dimeric naphthoquinone xanthomegnin and juglone at low concentrations uncoupled oxidative phosphorylation of intact rat liver mitochondria [129, 130]. The uncoupling action of naphthoquinones was determined by phenolic hydroxyl groups because the methylated analogues lost their uncoupling properties [130]. Interestingly, the most bacterially active naphthoquinones (viomellein and dihydroviomellein) contained more hydroxyl groups [97], which can be a reflection of their uncoupling activity.

Of the other metabolites with antibiotic activities, the mechanism of action was investigated for gunacin and 2-hydroxyjuglone. Gunacin synthesized by the fungus *Ustilago* sp. [15] features a high antibacterial activity against Gram-positive bacteria, fungi and mycoplasmas. It was found that gunacin inhibited the inclusion of ^{14}C -labelled thiamine and uracil into DNA and RNA and did not inhibit the induction of interferon in fibroblasts. Based on these results, the authors concluded that the mechanism of the cytotoxic action of the naphthoquinone is related to the disturbance of DNA synthesis [15].

The mechanism of the cytotoxic action of 2-hydroxyjuglone [131] a metabolite of the fungi *Pyricularia oryzae* [24], *Verticillium dahliae* [22] and others [26], is slightly different from that of gunacin. It was shown [131] that 2-hydroxyjuglone inhibited the respiration of fungal spheroplasts and mitochondria, suppressed the biosynthesis of RNA, proteins, lipids and transport of glucose. These results, in the opinion of the authors, indicate that 2-hydroxyjuglone has an inhibi-

tory effect on the basic biochemical processes and membrane systems of the cells [131].

Considering the fact that the metabolites of the naphthoquinone structure are potentially capable of redox conversions, we suggested and then showed [60, 113–115, 121, 132, 133] that their antibiotic and phytotoxic effects are due to the interaction with the oxidative systems of microbial and plant cells. We have investigated the naphthoquinones of the fungi *Fusarium decemcellulare* (fusarubin, anhydrofusarubin, javanicin, anhydrojavanicin, bostricoidin, nor-javanicin) and *Verticillium dahliae* (flaviolin and 2-hydroxyjuglone) [60, 113, 114, 121].

It has been found that the respiration of the Gram-positive bacteria, yeasts and fungi is noticeably activated by naphthoquinones after its suppression by cyanide [60, 113, 121]. These data indicate that naphthoquinones are capable of interacting with the redox systems of sensitive cells. Naphthoquinones accepted the reducing equivalents from the redox enzymes and transferred them directly to oxygen.

Studies of the subcellular components of bacterial, yeast and fungal cells showed that such systems are soluble cytoplasmic, flavin-containing NAD(P)H-dependent diaphorases [121]. Interestingly, respiration of Gram-negative bacteria was not activated by the pigments. However, naphthoquinones catalysed the oxidation of NAD(P)H by the cytoplasmic fraction of the Gram-negative bacteria [121]. This indicates that the resistance of the Gram-negative bacteria to the pigments is due to the absence of their transport across the cytoplasmic membrane.

In yeasts and fungi, the pigments actively catalysed the oxidation of the exogenous NAD(P)H by mitochondria in the presence of cyanide. It proved that the naphthoquinones accepted the reducing equivalents from flavin of the exogenous NADH dehydrogenase and transferred them to oxygen, bypassing the respiratory chain [113, 121].

We should also note that some of the naphthoquinones (2-hydroxyjuglone) exhibited an inhibitory effect on NADH:ferricyanide-reductase activity of complex I of the respiratory chain of rat liver mitochondria [133]. These data indicate that the pigments are capable of disturbing ATP synthesis in mitochondria at the oxidation of NAD-dependent substrates.

The mechanism of the phytotoxic action of the fungal naphthoquinones was investigated on pea seedlings [114, 115]. It was shown that, as in the case of eucaryotic organisms, the metabolites of the *Fusarium* (javanicin, fusarubin, anhydrofusarubin, bostricoidin) and *Verticillium* (2-hydroxyjuglone, flaviolin) fungi actively catalysed the oxidation of NADH and NADPH by the soluble, mitochondrial and microsomal fractions of the cells in the presence of oxygen. As a result of the transfer of the reducing equivalents by the naphthoquinone pigments from the flavine NAD(P)H-dependent dehydrogenases to

oxygen there occurred the formation of superoxide radicals [114, 115].

It is known [122, 134, 135] that the one-electron reduction of quinone compounds is accompanied by the formation of semiquinone radicals. The subsequent auto-oxidation of the quinone radicals in the presence of molecular oxygen is coupled with the formation of superoxide anion radicals, potentially toxic species. Thus, the one-electron reduction of quinones to semiquinones and the subsequent auto-oxidation of semiquinones to quinones leads to the formation of a large amount of superoxide anions. This redox process, known as the quinone "redox cycle", determines the oxidative stress caused by naphthoquinones [122].

The quinone "redox cycle" is catalysed by various flavine enzymes. In bacteria, these are soluble DT diaphorases [60]. In yeasts, fungi and plants the reduction of naphthoquinones was performed by cytoplasmic soluble NADH and NADPH diaphorases, microsomal flavine NAD(P)H-dependent dehydrogenases and mitochondrial exogenous NADH dehydrogenase localized at the outer side of the inner membrane [60, 113–115]. In animal cells the naphthoquinone "redox cycle" is maintained by NADPH cytochrome P-450 reductase, NADH cytochrome b-5 reductase and NADH ubiquinone oxidoreductase [122]. A relative ability of each of these enzymes to catalyse one-electron reduction is determined by the redox potential of naphthoquinone.

Besides the enzyme, naphthoquinones are capable of being directly reduced by intracellular electron carriers. Thus, for instance, menadione and other naphthoquinones react with the reduced glutathione and SH groups of membrane proteins. In this case, compounds are formed that preserve the ability to perform the "redox cycle" to form superoxide radicals and other active oxygen forms [135]. Spontaneous or enzymatic dismutation of superoxide radicals is accompanied by the formation of oxygen peroxide and molecular oxygen. Besides, the superoxide radicals in the presence of some metal ions can react with oxygen peroxide to form more reactive compounds—hydroxyl radicals and singlet oxygen [134, 136]. Thus, getting into the cell, as a result of the flavoprotein-catalysed or nonenzymatic "redox cycle", naphthoquinones rapidly lead to oxidative stress. Formation of active oxygen forms is accompanied by damage to the DNAs, proteins and membranes [122, 134–136].

Further studies showed that naphthoquinones were capable of inhibiting the activity of glutathione reductase [132]. This enzyme responsible for the maintenance of a high concentration of reduced glutathione in the cell is crucial in the protection against the "oxidative stress"—an increased concentration of superoxide radicals and other active oxygen forms. Moreover, it proved [137] that the superoxide-generating quinones (1,4- and 1,2-naphthoquinones) inhibit the activity of superoxide dismutase—an




Fig. 2. Proposed mechanism of cytotoxic action of the naphthoquinone metabolites of fungi. Naphthoquinones interact with mitochondria, microsomes and cytosolic proteins and function as free radical carriers. They augment the flow of electrons from NAD(P)H to molecular oxygen. This reaction is catalysed by flavoproteins and produces a free radical intermediate form pigment. As free radicals, these pigments, because of their high affinity and selective binding to nucleic acids, have the potential to be "site-specific free radicals" that bind to DNA or RNA, and either react directly or generate oxygen-dependent free radicals such as superoxide radical or hydroxyl radical to cause the damage associated with their cytotoxic actions.

enzyme responsible for the decrease in the concentration of superoxide radicals.

On the whole, we believe the following two aspects to be characteristic of the mechanism of the cytotoxic action of the auto-oxidative naphthoquinone fungal metabolites (Fig. 2).

Firstly, the disturbance of the energy metabolism due to the uncontrollable oxidation of NADH and NADPH and, thus, their removal from the oxidative phosphorylation system as potential sources of reducing equivalents for the respiratory chain. Second, the disturbance of the constructive metabolism determined by the formation of chemically active forms of reduced oxygen and free-radical forms of the pigments capable of interaction with DNA and RNA.

In a sense, the mechanism of action of naphthoquinones can be called universal. In nature, there is a great number of compounds (benzoquinones, naphthoquinones, etc.) which possess auto-oxidative properties and are capable of evoking the "oxidative stress" when they get into a living cell.

THE MECHANISM OF RESISTANCE OF THE FUNGI TO THEIR OWN METABOLITES

Resistance of microbial producers to their own metabolites possessing a high biological activity is described in detail in a review by Vining [138]. The insensitivity of the organisms-producers to the toxic metabolites is noted to be achieved in two ways. Firstly, by the change of the sensitivity of the target or its absence in the cells of the producer. Second, by the change of the toxicity of the metabolite by phosphorylation, acetylation etc. This section pre-

sents the results of our studies on the mechanism of the tolerance of the fungus *F. decemcellulare* to its own naphthoquinones.

Biosynthesis and accumulation of naphthoquinones in the culture medium is observed under conditions of fungal growth inhibition due to the decrease of pH of the medium to 3.5 and lower [60, 107]. Under these conditions, the pigments formed had no effect on the metabolism of the producer [60]. An increase in pH of the medium to 6.0 resulted in an effect of the pigment on the fungi, which was manifest in the increase of the cyanide-insensitive respiration due to the interaction of the auto-oxidative pigments with the redox systems of the cells [60]. As it was to be expected, under conditions optimal for growth (pH of the cultivation medium, 5.5–6.5) naphthoquinones inhibited fungal growth. In the presence of purified individual pigments the lag phase noticeably increased—up to 20–22 hr as against 6 hr in their absence [64]. In this period the respiration of the fungi was almost twice as active as compared with the control and by 40–60% insensitive to cyanide. Besides, the cells in the presence of the pigment have 3 times as high superoxide dismutase activity and 2 times as high catalase activity. The fact that these enzymes are responsible for the decrease of the level of the active oxygen forms in the cells in oxidative stress suggests that under these conditions naphthoquinones actively function as auto-oxidative acceptors of reducing equivalents [60, 121].

By the beginning of the active fungal growth (the end of the lag phase) the pigments added were transformed by 80–90% to the inactive metabolites [64]. According to the data of mass, IR and UV spectroscopy, the transformation was due to the demethylation of the 7-methoxy group of the quinone ring. In *O*-demethyl derivatives of naphthoquinones the ability to accept the reducing equivalents from the flavine dehydrogenases of the fungi and bacteria decreases [64]. The decrease in the acceptor properties is, probably, due to the decrease in their redox potential as it was noted for other naphthoquinones [139].

It should be pointed out that the decrease in the activity by way of changing the nature of the substituent in the quinone ring can be of crucial importance in the protection of the producer from the toxic effect of the metabolites during their biosynthesis. As it was already noted, the synthesis of naphthoquinones is coupled at a certain stage with the formation of nonmethylated derivatives [118], which are not toxic for the producer. Presumably, methylation (i.e. activation) of naphthoquinones is the last reaction in their biosynthesis and is performed at the stage of excretion.

CONCLUSION

Fungal naphthoquinones are typical representatives of the secondary metabolites and are synthesized under conditions of inhibition or total cess-

ation of fungal growth [108, 116]. As for many other secondary metabolites, their physiological role for producers is still a matter of dispute and there is no common view on this problem [140, 141]. Since naphthoquinones possess a broad-range biological activity and a universal mechanism of action, they can play a significant ecological role as protectors by providing a selective advantage for the producer in its survival in a natural ecosystem. In this sense they are largely similar to the phytoalexins of plants (some of which are also of naphthoquinone structure) whose protective function is generally accepted [142]. The secondary metabolites in fungi, as phytoalexins in plants, are induced by factors of biotic and abiotic origin including the foreign competitive organisms, toxic compounds etc. [56, 142]. In other words, formation of biologically active metabolites is a reaction to unfavourable factors in the environment.

Most naphthoquinones are synthesized by phytopathogenic fungi (Table 2). At first glance, one should have expected that they shall play a significant role in the pathogenicity of the fungi with respect to plants. However, no direct evidence of this has been obtained as of today. Nevertheless, the compounds can actively change the metabolism of plant cells by suppressing their protection mechanisms.

REFERENCES

- Rice, E. L., *Botanical Reviews*, 1979, **45**, 15.
- Thomson, R. H., *Naturally Occurring Quinones*. Academic Press, London, 1971, p. 198.
- Parisot, D., *Microbiosystems*, 1990, **64**, 31.
- Tanaka, H., Koyama, Y., Awaya, J., Marumo, H., Oiwa, R., Katagiri, M., Nagai, T. and Omura, S., *Journal of Antibiotics*, 1975, **28**, 860.
- Kern, H., *Annals Phytopathology*, 1978, **10**, 327.
- Baker, R. A., Tatum, J. H. and Nemeč, S., Jr., *Phytopathology*, 1981, **71**, 951.
- Marcinkowska, J., Kraft, J. M. and Marquis, L. M., *Canadian Journal of Plant Science*, 1982, **62**, 1027.
- Nemeč, S., Baker, R. A. and Tatum, J. H., *Soil Biology and Biochemistry*, 1988, **20**, 493.
- Claydon, N., Grove, J. F. and Pople, M., *Journal of Invertebrate Pathology*, 1977, **30**, 216.
- Baker, R. A., Tatum, J. H. and Nemeč, S. J., *Mycopatologia*, 1990, **111**, 9.
- Arnstein, H. R. and Cook, A. H., *Journal of the Chemical Society*, 1947, 1021.
- Kern, H. and Naef-Roth, S., *Phytopathologie Zeitschrift*, 1967, **60**, 316.
- Kurobane, I., Zaita, N. and Fukuda, A., *Journal of Antibiotics*, 1986, **39**, 205.
- Tatum, J. H., Baker, R. A. and Berry, R. E., *Phytochemistry*, 1987, **26**, 2499.
- Werner, R. G., Appel, K.-R. and Merk, W. M. A., *Journal of Antibiotics*, 1979, **32**, 1104.
- Tanaka, H., Marumo, H., Nagai, T., Okada, M., Taniguchi, K. and Omura, S., *Journal of Antibiotics*, 1975, **28**, 925.
- Balan, J., Fuska, J., Kuhr, I. and Kuhrova, V., *Folia Microbiologica*, 1970, **15**, 479.
- Vogel, F. S., Kemper, L. A. K., McGarry, S. J. and Graham, D. G., *American Journal of Pathology*, 1975, **78**, 33.
- Lown, J. W. and Sim, S. K., *Canadian Journal of Biochemistry*, 1976, **54**, 446.
- Shucla, J. N., Tandon, J. S., Bharuni, D. S. and Dhar, M. M., *Phytochemistry*, 1971, **10**, 1909.
- Bendz, G., *Acta Chemica Scandinavica*, 1948, **2**, 192.
- Stipanovic, R. D. and Bell, A. A., *Mycologia*, 1977, **69**, 164.
- Soga, O., *Zeitschrift für Naturforschung*, 1976, **31**, 124.
- Yamagushi, I., Sekido, S. and Misato, T., *Journal of Pesticide Science*, 1983, **8**, 229.
- Bell, A. A., Stipanovic, R. D. and Puhalla, J. E., *Tetrahedron*, 1976, **32**, 1353.
- Zhdanova, N. N., Stepanichenko, N. I., Vasilevskaya, A. I., Navrezova, N. S., Tyschenko, A. P., Mukchamedzhanov, S. Z. and Aslanov, Kh. A., *Microbiologichny Journal*, 1985, **47**, 43.
- McGovern, E. P. and Bentley, M. R., *Biochemistry*, 1975, **14**, 3314.
- Astil, B. D. and Roberts, J. C., *Journal of the Chemical Society*, 1953, 3303.
- Knapp, J. E., Chao, P.-D., Shiff, P. L. and Slatkin, D. J., *Journal of Chemical Research*, 1979, 2685.
- Assante, G., Locci, R., Camarda, L., Merlini, L. and Nasini, G., *Phytochemistry*, 1977, **16**, 243.
- Soga, O. and Iwamoto, H., *Zeitschrift für Naturforschung*, 1980, **35**, 1497.
- Chimura, H., Sawa, T., Kumada, Y., Nakamura, F., Matsusaki, M., Takita, T., Takeuchi, T. and Umezawa, H., *Journal of Antibiotics*, 1973, **26**, 618.
- Howe, R. and Moore, R. H., *Experientia*, 1969, **25**, 474.
- Van Eijk, J. W. and Roemans, H. J., *Experientia*, 1978, **34**, 1257.
- Natori, S., Kumada, Y. and Nishikawa, H., *Chemical and Pharmaceutical Bulletin (Tokyo)*, 1965, **13**, 633.
- Bentley, R. and Gatenbeck, S., *Biochemistry*, 1965, **4**, 1150.
- Kurobane, I., Vining, L. C., McInnes, A. G. and Walter, J. A., *Canadian Journal of Chemistry*, 1980, **58**, 1380.
- Van der Kerk, G. J. M. and Overeem, J. C., *Recueil des travaux chimiques des Pays-Bas*, 1964, **83**, 995.
- Kobayashi, A., Yata, S., Nino, T. and Kawasu, K., *Agricultural and Biological Chemistry*, 1987, **51**, 2857.
- Kopanski, L., Besl, H. and Steglich, W., *Pure and Applied Chemistry*, 1981, **53**, 1233.

41. Coombe, R. G., Lowe, H. I. C. and Watson, T. R., *Australian Journal of Chemistry*, 1972, **25**, 875.
42. Krivoshchekova, O. E., Maximov, O. B., Stepanenko, L. S. and Mishchenko, N. P., *Phytochemistry*, 1982, **21**, 193.
43. Brunn, T. and Lamvik, A., *Acta Chemica Scandinavica*, 1971, **25**, 483.
44. Kadkol, M. V., Gopalkrishnan, K. S. and Narasimhachari, N., *Journal of Antibiotics*, 1971, **24**, 245.
45. Narasimhachari, N. and Gopalkrishnan, K. S., *Journal of Antibiotics*, 1974, **27**, 283.
46. Armstrong, J. J. and Turner, W. B., *Journal of the Chemical Society*, 1965, 5927.
47. Van Eijk, G. W. and Roeymans, H. J., *Phytochemistry*, 1978, **17**, 1804.
48. Closs, A. and Sigg, H. P., *Helvetica Chimica Acta*, 1973, **56**, 619.
49. Baker, P. M. and Bullock, E., *Canadian Journal of Chemistry*, 1969, **47**, 2733.
50. Huneck, S., Steglich, W. and Hofle, G., *Phytochemistry*, 1977, **16**, 121.
51. Donnelly, D. and O'Reilly, J., *Phytochemistry*, 1980, **19**, 277.
52. Birch, A. J., Butler, D. N., Effenberger, R., Rickards, R. W. and Simpson, T. G., *Journal of the Chemical Society, Perkin Transactions I*, 1979, 807.
53. Effenberger, R. and Simpson, T. J., *Journal of the Chemical Society, Perkin Transactions I*, 1979, 823.
54. Cross, B. E., Edinberry, M. N. and Turner, W. D., *Chemical Communications*, 1970, 209.
55. Cross, B. E. and Zammitt, L. J., *Journal of the Chemical Society, Perkin Transactions I*, 1973, 2975.
56. Parisot, D., Maugin, M. and Gerlinger, C., *Journal of Genetics and Microbiology*, 1981, **126**, 443.
57. Roos, A., *Phytopathologie Zeitschrift*, 1977, **88**, 238.
58. Ruelius, H. W. and Gauhe, A., *Liebigs Annalen der Chemie*, 1950, **568**, 38.
59. Natori, S., Inouye, Y. and Nishikawa, H., *Chemical and Pharmaceutical Bulletin (Tokyo)*, 1967, **15**, 380.
60. Medentsev, A. G., Baskunov, B. P. and Akimenko, V. K., *Biokhimiya*, 1988, **53**, 413.
61. Steyn, P. S., Wessels, P. L. and Marasas, W. F. O., *Tetrahedron*, 1979, **35**, 1551.
62. Parisot, D., Devys, M. and Barbier, M., *Phytochemistry*, 1985, **24**, 1977.
63. Tatum, J. H., Baker, J. H. and Berry, R. E., *Phytochemistry*, 1985, **24**, 457.
64. Medentsev, A. G., Baskunov, B. P. and Akimenko, V. K., *Biokhimiya*, 1989, **54**, 619.
65. Tatum, J. H. and Baker, R. A., *Phytochemistry*, 1983, **22**, 543.
66. Gerber, N. N. and Ammar, M. S., *Journal of Antibiotics*, 1979, **32**, 685.
67. Kurobane, I., Vining, L. C., McInnes, A. G. and Smith, D. G., *Canadian Journal of Chemistry*, 1978, **56**, 1593.
68. Barbier, M., Devys, M. and Parisot, D., *Canadian Journal of Chemistry*, 1988, **66**, 2803.
69. Parisot, D., Devys, M. and Barbier, M., *Journal of Antibiotics*, 1991, **44**, 103.
70. Parisot, D., Maugin, M. and Gerlinger, C., *Journal of Genetics and Microbiology*, 1984, **130**, 1543.
71. Tatum, J. H., Baker, R. A. and Berry, R. E., *Phytochemistry*, 1989, **28**, 283.
72. Parisot, D., Ferezou, J.-P. and Barbier, M., *Phytochemistry*, 1983, **22**, 1301.
73. Parisot, D., Devys, M. and Barbier, M., *Phytochemistry*, 1989, **28**, 3240.
74. Cross, B. E., Myers, P. L. and Webster, G. R. B., *Journal of the Chemical Society*, 1970, 930.
75. Kimura, Y., Hamasaki, T. and Nakajima, H., *Agricultural and Biological Chemistry*, 1981, **45**, 2653.
76. Kimura, Y., Shimada, A., Hamasaki, T. and Nakajima, H., *Agricultural Biological Chemistry*, 1988, **52**, 1253.
77. Tatum, J. H., Baker, R. A. and Berry, R. E., *Phytochemistry*, 1985, **24**, 3019.
78. Shilton, W. S., *Journal of Organic Chemistry*, 1968, **33**, 4295.
79. Parisot, D., Devys, M. and Barbier, M., *Microbios Letters*, 1987, **36**, 129.
80. Arsenault, G. P., *Tetrahedron*, 1968, **24**, 4745.
81. Cajori, F. A., Otani, T. T. and Hamilton, M. A., *Journal of Biological Chemistry*, 1954, **208**, 107.
82. Kurobane, I., Vining, L. C., McInnes, A. G. and Gerber, N. N., *Journal of Antibiotics*, 1980, **33**, 1376.
83. Parisot, D., Devys, M. and Barbier, M., *Zeitschrift für Naturforschung*, 1989, **44B**, 1473.
84. Parisot, D., Devys, M. and Barbier, M., *Phytochemistry*, 1990, **29**, 3364.
85. Parisot, D., Devys, M. and Barbier, M., *Phytochemistry*, 1988, **27**, 3002.
86. Medentsev, A. G., Fayn, M. E., Aytchodzina, N. A. and Nikitina, E. T., *Vestnik Akademii Nauk Kazachstana*, 1991, **12**, 70.
87. Guy Valadon, L. R. and Chapman, D. J., *Microbios Letters*, 1983, **24**, 115.
88. Cornforth, J. W., Ryback, G. and Robinson, P. M., *Journal of the Chemical Society*, 1971, 2786.
89. Kjaer, D., Kjaer, A., Pedersen, C., Bu'Lock, J. D. and Smith, J. R., *Journal of the Chemical Society*, 1971, 2792.
90. Stipanovic, R. D. and Wheeler, M. H., *Pesticide and Biochemical Physiology*, 1980, **13**, 198.
91. Wirth, J. G., Beesley, T. E. and Anand, S. R., *Phytochemistry*, 1965, **24**, 505.
92. Blank, F., Day, W. C. and Just, G., *Journal of Investigative Dermatology*, 1963, **40**, 133.
93. Ng, A. S., Just, G. and Blank, F., *Canadian Journal of Chemistry*, 1969, **47**, 1223.

94. Durley, R. C., McMillan, J., Simpson, T. J., Glen, A. T. and Turner, W. D., *Journal of the Chemical Society, Perkin Transactions I*, 1975, 163.
95. Stack, M. E., Eppley, R. M., Dreifuss, P. A. and Pohland, A. E., *Applied Environmental Microbiology*, 1977, **33**, 351.
96. Nozawa, Y. and Ito, Y., *Experientia*, 1970, **26**, 803.
97. Sedmera, P., Volk, J., Weijer, J., Vokoun, J. and Musilek, V., *Collection of Czechoslovak Chemical Communications*, 1981, **46**, 1210.
98. Zeeck, A., Russ, P., Laatsch, H., Loeffler, W., Wehrle, H. and Zahner, H., *Chemische Bericht*, 1979, **112**, 957.
99. Stack, M. E., Mazzola, E. P. and Eppley, R. M., *Tetrahedron*, 1979, **52**, 4989.
100. Akita, T., Kawai, K., Shimonaka, H., Nozawa, Y., Yto, Y. and Nishube, S., *Shinkin to Shikinisho*, 1975, **16**, 177.
101. Birchall, G. R., Bowden, K., Weiss, U. and Whalley, W. D., *Journal of the Chemical Society*, 1966, 2237.
102. Ashley, J. W., Hobbs, B. C. and Raistrick, H., *Biochemical Journal*, 1937, **31**, 385.
103. Shibata, S., Morshita, E., Takeda, T. and Sakata, K., *Chemical and Pharmaceutical Bulletin (Tokyo)*, 1968, **16**, 405.
104. Medentsev, A. G., Kotik, A. N., Truphanova, V. A. and Akimenko, V. K., *Prikladnaya Biokhimiya i Mikrobiologiya*, 1993, **29**, 542.
105. Baker, P. M. and Roberts, J. C., *Journal of the Chemical Society*, 1966, 2234.
106. Takeda, T., Morishita, E. and Shibata, S., *Chemical and Pharmaceutical Bulletin (Tokyo)*, 1968, **16**, 2213.
107. Medentsev, A. G. and Akimenko, V. K., *Mikrobiologiya*, 1992, **61**, 824.
108. King, T. S., Roberts, J. C. and Thomson, D. J., *Chemical Communications*, 1970, 1499.
109. Bleckburn, G. M., Neilson, A. N. and Todd, L., *Proceedings of the Chemical Society*, 1962, 327.
110. Read, G., Rashid, A. and Vining, L. C., *Journal of the Chemical Society*, 1969, 2059.
111. Gill, M. and Gimenez, A., *Journal of the Chemical Society, Perkin Transactions I*, 1995, 645.
112. Kern, H., Naef-Roth, S. and Ruffner, F., *Phytopathologie Zeitschrift*, 1972, **74**, 272.
113. Medentsev, A. G. and Akimenko, V. K., *Biokhimiya*, 1989, **54**, 1904.
114. Medentsev, G., Maslov, A. N. and Akimenko, V. K., *Biokhimiya*, 1990, **55**, 1766.
115. Medentsev, A. G. and Akimenko, V. K., *Phytochemistry*, 1992, **31**, 77.
116. Bu'Lock, J. D., *The Biosynthesis of Mycotoxins, A Study in Secondary Metabolism*. Academic Press, New York, 1980, p. 1.
117. Holenstein, J. E., Stoessl, A., Kern, H. A. and Stothers, J. B., *Canadian Journal of Chemistry*, 1984, **62**, 1971.
118. Gatenbeck, S. and Bentley, R., *Biochemical Journal*, 1965, **94**, 478.
119. Arnstein, V. R., Cook, A. H. and Lacey, M. S., *Nature*, 1946, **157**, 333.
120. Ammar, M. S., Gerber, N. N. and McDaniel, L. E., *Journal of Antibiotics*, 1979, **32**, 679.
121. Medentsev, A. G. and Akimenko, V. K., *Biokhimiya*, 1988, **53**, 289.
122. Smith, M. T., Evans, C. G., Thor, H. and Orrenius, S., in *Oxidative Stress*. Academic Press, London, 1985, p. 91.
123. Carlton, W. W., Stack, M. E. and Eppley, R. M., *Toxicology and Applied Pharmacology*, 1976, **38**, 455.
124. Kern, H. and Naef-Roth, S., *Phytopathologie Zeitschrift*, 1965, **53**, 45.
125. Kern, H., Naef-Roth, S. and Item, H., *Phytopathologie Zeitschrift*, 1970, **67**, 1.
126. Kern, H., in *Phytotoxins in Plant Disease*. Academic Press, London, 1972, p. 35.
127. Kern, H. and Naef-Roth, S., *Phytopathologie Zeitschrift*, 1966, **57**, 289.
128. Dorn, S., *Phytopathologie Zeitschrift*, 1974, **81**, 193.
129. Ito, Y., Kawai, K. and Nozawa, Y., *Journal of Biochemistry*, 1973, **74**, 805.
130. Kawai, K., Akita, T., Nishibe, S., Nozawa, Y., Ogihara, Y. and Yto, Y., *Journal of Biochemistry*, 1976, **79**, 145.
131. Yamagushi, I., Sekido, S., Seto, H. and Misato, T., *Journal of Pesticide Science*, 1983, **8**, 545.
132. Bironaite, D. A., Cenas, N. K., Medentsev, A. G., Kulis, J. J. and Akimenko, V. K., *Zeitschrift für Naturforschung*, 1991, **46c**, 966.
133. Bironaite, D. A., Cenas, N. K., Anusevicius, Z. J., Medentsev, A. G. and Akimenko, V. K., *Archiv der Biochimica et Biophysica*, 1992, **297**, 253.
134. Emanuel, N. M., Bogdanov, G. I. and Orlov, V. S., *Uspekhi Khimii*, 1984, **53**, 1929.
135. Wefers, H. and Sies, H., *Archiv der Biochimica et Biophysica*, 1983, **224**, 568.
136. Kappus, H. and Sies, H., *Experientia*, 1981, **37**, 1233.
137. Smith, M. T. and Evans, C. G., *Biochemical Pharmacology*, 1984, **33**, 3109.
138. Vining, L. C., *Advances in Applied Microbiology*, 1979, **25**, 147.
139. Hodnett, E. M., Wongweichintana, C., Dunn, III, W. J. and Marrs, P., *Journal of Medical Chemistry*, 1983, **26**, 570.
140. Vinning, L. C., *Biotechnology*, 1986, **4**, 20.
141. Martin, J. F. and Demain, A. L., *Microbiological Review*, 1980, **44**, 230.
142. Darvill, A. G. and Albersheim, P., *Annual Review of Plant Physiology*, 1984, **35**, 243.