

Vera Meyer · Ulf Stahl

New insights in the regulation of the *afp* gene encoding the antifungal protein of *Aspergillus giganteus*

Published online: 5 July 2005
© Springer-Verlag 2005

Curr Genet (2002) 42:36–42

In the issue 42 of the Journal Current Genetics, in the article entitled “New insights in the regulation of the *afp* gene encoding the antifungal protein of *Aspergillus giganteus*” by Meyer and Stahl (pp 36-42), the sequence and positions of the two putative PacC binding sites *afpP1* and *afpP2* were mistakenly interchanged in the text and in Table 1. Thus, the following changes should be made in the text:

- Page 36, right column, lines 2–3 up: “*afpP1* (3'-GAACCG-5') and *afpP2* (5'-GCCAAG-3')” should be “*afpP1* (5'-GCCAAG-3') and *afpP2* (3'-GAACCG-5')”
- Page 38, left column, lines 12–13 up: “from –937 to –932 and from –1138 to –1133” should be “from –1138 to –1133 and from –937 to –932” In Table 1, name “*afpP1*” should be replaced with “*afpP2*” and vice versa. The corrected Table 1 is reprinted below.
The authors wish to apologize for the error.

Table 1 Oligonucleotides used in this study. The position of each oligonucleotide in the promoter and coding region of the *afp* gene is given. Numbering is from the first codon of translation. The nucleotides GCG AAT TCA at the 5' end of EI do not correspond to the *afp* promoter. Putative PacC binding sites are shown in *italics*

Primer	Sequence	Source	Name	Position
Pac20	5'-TGG TTT A <i>GC CAA GGG</i> GAA CG-3'	<i>afp</i> promoter	<i>afpP1</i>	From –1145 to –1126
Pac(–)20	5'-CGT TCC C <i>CTTGGCTA</i> AAC CA-3'	<i>afp</i> promoter	<i>afpP1</i>	From –1126 to –1145
Pac30	5'-GAG CGC ATA TGA AGT <i>ACCTTGCG</i> GCT TAG-3'	<i>afp</i> promoter	<i>afpP2</i>	From –956 to –927
Pac(–)30	5'-AGC TCT AAG CC <i>G CCA</i> AGG TAC TTC ATA TGC-3'	<i>afp</i> promoter	<i>afpP2</i>	From –923 to –952
Pac22	5'-GAT GAT CAG CCA ACT GAA AAC A-3'	<i>afp</i> promoter	<i>afpP3</i>	From –1120 to –1099
Pac(–)22	5'-TGT TTT CAG TTG GCT GAT CAT C-3'	<i>afp</i> promoter	<i>afpP3</i>	From –1099 to –1120
EI	5'-GCG AAT TCA CAG TTA CGG ACA ATC GAT TG-3'	<i>afp</i> promoter	–	From –1100 to –980
Pro10(–)	5'-TCT CTG CAA TGA ACA AGG AAC-3'	<i>afp</i> promoter	–	From –884 to –904
Pac18	5'-GCT GCC <i>CTTGGC</i> GCG GTT-3'	<i>afp</i> gene	<i>afpP4</i>	From 43 to 60
Pac(–)18	5'-AAC CGC <i>GCCAAG</i> GGC AGC-3'	<i>afp</i> gene	<i>afpP4</i>	From 60 to 43
OlantiA	5'-GCC ACA TAC AAT GGC GTA GGT-3'	<i>afp</i> gene	OIA	From 130 to 150
Olanti(–)A	5'-ACC TAC GCC ATT GTA TGT GGC-3'	<i>afp</i> gene	OIA	From 150 to 130
OlantiE	5'-CTA GCA GTA GCA CTT CCC CTT-3'	<i>afp</i> gene	–	From 432 to 412
AnidPac1	5'-GGG GCG GCT GCG ACA GCT GCT GTG G-3'	<i>pacC</i> gene, <i>Aspergillus nidulans</i>	–	–
AnidPac2	5'-AGG CAT TTT GGA GTC TCC ATC CTC ATC-3'	<i>pacC</i> gene, <i>A. nidulans</i>	–	–

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00294-002-0324-9>

V. Meyer (✉) · U. Stahl
Technische Universität Berlin, Institut für Biotechnologie,
Fachgebiet Mikrobiologie und Genetik,
Gustav-Meyer-Allee 25, 13355 Berlin, Germany
E-mail: V.Meyer@LB.TU-Berlin.de