



ELSEVIER

Food Chemistry 65 (1999) 139–151

**Food
Chemistry**

Review

2,5-Dimethyl-4-hydroxy-2*H*-furan-3-one and its derivatives: analysis, synthesis and biosynthesis—a review

I. Zabetakis*, J.W. Gramshaw, D.S. Robinson

Procter Department of Food Science, The University of Leeds, Leeds, LS2 9JT, UK

Received 18 May 1998; received in revised form and accepted 14 September 1998

Abstract

2,5-Dimethyl-4-hydroxy-2*H*-furan-3-one is believed to be a key flavour constituent in many fruits and baked foods. The analytical and organic methods applied to the analysis of DMHF and its derivatives, synthesis and biosynthesis are reviewed. Possibilities for further elucidation of biosynthetic pathways and the biotechnological production of DMHF are considered. © 1999 Elsevier Science Ltd. All rights reserved.

1. The analysis of 2,5-dimethyl-4-hydroxy-2*H*-furan-3-one and its derivatives

2,5-Dimethyl-4-hydroxy-2*H*-furan-3-one (DMHF) is widely distributed in nature and has a very low flavour threshold value in water (4×10^{-5} mg/kg); hence, its effect on food aroma is considerable (Latrasse, 1991). 2,5-Dimethyl-4-hydroxy-2*H*-furan-3-one occurs in nature in four forms (Fig. 1): 2,5-dimethyl-4-hydroxy-2*H*-furan-3-one glucoside (DMHF glucoside), 2,5-dimethyl-4-hydroxy-2*H*-furan-3-one 6'-*O*-malonyl- β -D-glucopyranoside (DMHF malonyl-glucoside), 2,5-dimethyl-4-methoxy-2*H*-furan-3-one (mesifuran) as well as the free aglycone (Latrasse, 1991; Mayerl, Naf, & Thomas, 1989; Roscher, Herderich, Steffen, Schreier, & Schwab, 1996a).

Recently, DMHF has attracted the interest of many researchers working in the field of flavour analysis (Pickenhagen, Velluz, Passerat, & Ohloff, 1981), aroma generation through chemical (Schieberle, 1992) and enzymatic processes (Roscher, Schreier, & Schwab, 1997c), organic synthesis using enzymes as catalysts (Wong & Whitesides, 1985) and the sensory evaluation of complex flavour mixtures (Larsen, Poll, & Olsen, 1992). Analytical techniques such as HPLC with either UV (Lee & Nagy, 1987) or Photo Diode Array (PDA) detection (Zabetakis, Gramshaw, & Robinson, 1996), GLC-FID (Williams & Mottram, 1981) and GLC-MS

(Withopf, Richling, Roscher, Schwab, & Schreier, 1997) have been used for the analysis of DMHF and derivatives. Recently, the first studies on the biosynthesis of DMHF have taken place using either strawberry fruits (Hong, Huang, Reineccius, Harlander, & Labuza, 1990; Roscher et al., 1997c) or callus cultures (Zabetakis & Holden, 1996). Plant tissue culture techniques may provide useful tools for the elucidation of the biosynthetic pathway of strawberry flavour molecules, including manipulation of culture conditions to increase the net yield and further genetic manipulation (Zabetakis & Holden, 1997).

In this review are considered first the analysis of sources where the furanones occur naturally (e.g. fruits and other unprocessed material) and, second foodstuffs where occurrence results from exogenous processes, including the baking of bread, cooking of meat and manufacture of preserves.

1.1. Naturally occurring furanones

Free DMHF was the first furanone isolated and identified, simultaneously in pineapple (Rodin, Himel, Silverstein, Leeper, & Gortner, 1965) and strawberry (Willhalm, Stoll, & Thomas, 1965). The aroma of DMHF has been described as having a rotten note (Rodin et al.), a caramel-like note (Shaw, Tatum, & Berry, 1969; Tonsbeek, Plancken, & Weerdhof, 1968), as being sweet, floral and fruity (Miller, Libbey, & Yang, 1973) and becoming fruity and strawberry-like at low concentrations (Re, Mauren, & Ohloff, 1973).

* Corresponding author. E-mail: y.zabetakis@food.leeds.ac.uk




Fig. 1. The structures of 2,5-dimethyl-4-hydroxy-2*H*-furan-3-one, 2,5-dimethyl-4-methoxy-2*H*-furan-3-one, 2,5-dimethyl-4-hydroxy-2*H*-furan-3-one glucoside and 2,5-dimethyl-4-hydroxy-2*H*-furan-3-one malonyl-glucoside.

Later, DMHF glucoside was identified, firstly in strawberry (Mayerl et al., 1989); since then it has been regarded as the precursor of DMHF in all the fresh sources (eg fruits) where both compounds exist. It has also been detected in pineapple (Wu et al., 1990) and tomatoes (Krammer, Takeoka, & Buttery, 1994). The amounts of DMHF glucoside in pineapple (Wu, Kuo, Hartman, Rosen, & Ho, 1991) and strawberry (Wintoch, Krammer, & Schreier, 1991; Sanz, Perez, & Richardson, 1994) have been measured (Table 1). Recently, 2,5-dimethyl-4-hydroxy-2*H*-furan-3-one 6'-O-malonyl- β -D-glucopyranoside (DMHF malonyl-glucoside) has been identified in strawberry fruit (*Fragaria* \times *ananassa*, cv. Senga Sengana) (Roscher et al., 1996a) with a ratio of the malonylated glucoside to its non-acylated glucoconjugate of 1:1 in the ripe fruit. Its presence suggests that the glucosides may be important derivatives of free DMHF and that their biosynthesis is therefore worthy of further study.

The synthesis of phenyl 6'-O-malonyl- β -D-glucopyranoside has been reported (Roscher, Steffen, Herderich, Schwab, & Schreier, 1996b). This compound together with the synthesized 6'-O-malonylated- β -D-glucopyranosides of benzyl alcohol, 2-phenylethanol, geraniol, citronellol and DMHF have been used as references for the spectroscopic identification of malonylated glucosides in plants (Withopf et al., 1997). This report indicated that malonylation of the glucoconjugates may be a common pathway in plant secondary metabolism. DMHF malonyl-glucoside was identified in raspberry and strawberry.

Table 1

The amounts of DMHF, DMHF glucoside and mesifuran in some fruits (mg of analyte kg^{-1} of fruit)

Fruit	DMHF	DMHF glucoside	Mesifuran
Strawberry (cv. Elsanta) ^a	20.9	4.3	79.5
Wild strawberry ^a	16.7	10.9	12.1
Wild raspberries ^b	0.1	nd ^j	tr ^j
Arctic bramble ^c	nd ^j	nd ^j	18
Alphonso mango ^d	2	nd ^j	nd ^j
Pineapple (from Ivory Coast) ^e	7.4	nd ^j	0.2
Pineapple (from Costa Rica) ^f	0.7	0.4	nd ^j
Tomato (cv. Ace) ^g	0.7	nd ^j	nd ^j
Tomato (var. 6203) ^h	nd ^j	tr ^j	nd ^j
Wine (various grape cultivars) ⁱ	0.1–3.5	nd	nd

^a Zabetakis and Holden, 1995.

^b Honkanen et al., 1980.

^c Kallio, 1976b.

^d Idstein and Schreier, 1985.

^e Pickenhagen et al., 1981.

^f Wu et al., 1991.

^g Krammer et al., 1994.

^h Buttery et al., 1995.

ⁱ Guedes de Pinho and Bertrand, 1995.

^j tr: traces, nd: not detected.

Mesifuran, described elsewhere as having a Xeres wine-like note (Hunter, Busek, & Radford, 1974), has been claimed to be the major flavour-active component of the arctic bramble (Pyysalo, Suihko, & Honkanen, 1977). The odour value of mesifuran was estimated to be extremely high (600,000), whereas the odour value

reported for DMHF was less than 100. Previously it was claimed, for arctic bramble, that the contribution of mesifuran to the total volatiles increased from 0.2 to 20% of the volatiles during ripening in the field (Kallio, 1976a).

Many of these furanones occur as a racemate in natural products (strawberry, pineapple, grape and wine) (Mosandl, Bruche, Askari, & Schmarr, 1990) and therefore stereospecific analysis is not useful for assessment of the authenticity of these commodities, or their derived products (Brueche et al., 1991). DMHF was identified in fresh and processed tomato and its contribution to aroma assessed (Buttery, Takeoka, Kramer, & Ling, 1994). Calculations of the concentration/threshold ratios have shown that DMHF occurs at a concentration well above its threshold and it is, thus, among the 10 compounds with the highest probability of contributing to both fresh and processed tomato flavour (Buttery, Takeoka, & Ling, 1995).

2,5-Dimethyl-4-hydroxy-2*H*-furan-3-one was also identified as one of 54 constituents of chempedak fruit (Wong, Lim, & Wong, 1992). A commercial lovage (*Levisticum officinale* Koch.) extract was found to contain six compounds of high sensory relevance, one of them being DMHF (Blank & Schieberle, 1993). Both DMHF and mesifuran were identified in the Finnish wild raspberry by Honkanen, Pyysalo, and Hirvi (1980) using GC–MS. In this study, more than 40 compounds not reported previously as raspberry volatiles were detected and included 11 terpenes as well as DMHF and mesifuran. Studies on mango aroma by high resolution GC/MS have revealed that it contained at least 152 aroma compounds and that DMHF is a major constituent (Idstein & Schreier, 1985). However, when the importance of some lactones and DMHF to mango aroma was assessed by Wilson, Shaw, and Knight (1990), it was claimed that DMHF did not make a positive contribution to the flavour.

A comparison of the amounts of DMHF and mesifuran in strawberries, pineapples and mangoes was carried out by GC after ethyl ether extraction (Pickenhagen et al., 1981). The contribution of DMHF to pineapple flavour was also reported in a review of the flavour of this fruit by Karg (1983), who claimed that the two essential flavour compounds of pineapple flavour were DMHF and 3-methyl mercaptopropionic acid. In a more recent study of pineapple flavour using capillary GC coupled to Fourier transform IR spectroscopy, Fehl and Marcott (1989) identified DMHF and phenylacetaldehyde, an unwanted off-flavour compound. A convenient reversed-phase HPLC method for the analysis of DMHF in pineapple and grapefruit juices was developed by Lee and Nagy (1987). This method is fast and convenient and avoids heating problems associated with the GC analysis of unstable compounds, such as DMHF (Williams & Mottram, 1981).

Use of GC–MS to identify DMHF and mesifuran (compounds responsible for a strawberry-like flavour

note) together with nasal appraisal by sniffing the GC eluate led to the identification of these furanones for the first time in berries and wines of interspecific grapevine breeding (Rapp, Knipser, & Engel, 1980). This report indicates that an objective diagnosis method could be developed in order to recognize the furanones at the seedling stage. Rapp, Knipser, Engel, and Hastrich (1983) have identified DMHF, in various grape varieties, as the compound responsible for a strawberry aroma which can be troublesome for some *Vitis vinifera* × *labrusca* (European × American) hybrids. In a comparative study of various cultivars of grapes, mesifuran was claimed to be responsible for the caramel-like undertone in the aroma fraction obtained from Concord and Niagara grapes, where it was detected in substantial amounts (0.15 and 0.18 mg/kg, respectively) but only found in trace amounts in Elvira (*V. labrusca*, L. × *V. riparia*, M.) grapes (Schreier & Paroschy, 1981).

However, analysis of the red grape juice of the Isabella (*V. vinifera* × *V. labrusca*) cultivar showed that this juice contained all the characteristic compounds of *V. labrusca* grapes, except for DMHF (Cesare & Nani, 1992). A recent paper describes the analytical determination of DMHF in wines by GC–MS and the possible application of this analysis to differentiate white wines from hybrids and various *V. vinifera* cultivars (Guedes de Pinho & Bertrand, 1995). Sixty-five wines of various hybrids and more than an hundred wines have been analysed using this method. It has been reported that vinification with grape skin contact leads to a decrease in DMHF levels and suggests that DMHF concentration can be increased by using pectolytic enzymes with -glucosidic secondary activities. In a study on the aroma of strawberry wine, Schreier and Drawert (1981) proposed that concentrations of fruit-derived aroma compounds such as mesifuran, methyl cinnamate and β -decalactone could be measured and used as a reference in order to detect adulteration of wines with aroma compounds.

Larsen et al. (1992), who considered the concentrations of aroma components in the light of threshold values, found DMHF, together with linalool and ethyl hexanoate to be important strawberry aroma compounds. In a complementary study, the odour thresholds of 24 aroma compounds in the strawberry cultivar Senga Sengana were determined using duo-trio tests; DMHF was found to be one of the 3 most important compounds (Larsen & Poll, 1992). Since, in most of the reports, the analysis of DMHF and derivatives was carried out by GC, a very useful study of the gas chromatographic analysis of DMHF has been carried out by Williams and Mottram (1981). They compared five types of capillary columns and illustrated the advantage of fused silica capillary columns for the analysis of potentially unstable compounds, such as DMHF.

Characterisation of the aroma of six strawberry cultivars and quantification of DMHF and mesifuran has

been reported by Douillard and Guichard (1990). The volatiles of the strawberries were isolated by direct solvent extraction using dichloromethane. Sixty compounds were identified and quantified by GC–MS. In this study, hept-2-en-3-ol was used as the internal standard for further analysis by GC–MS. The recovery factors of the analytes were assumed, however, to be equal to that of the internal standard—which could lead to incorrect quantitative estimations and partly explain the quite different results obtained by these authors when compared to those of Larsen et al. (1992). Supercritical Fluid Extraction (SFE) has been studied as a preparative tool for strawberry aroma analysis and compared to traditional solvent extraction using dichloromethane (Polesello, Lovati, Rizzolo, & Rovida, 1993). It was found that, using SFE, it was possible to recover the majority of the aroma compounds including the character impact compound of wild strawberries (mesifuran). However, this extraction method was more selective (i.e. fewer aromatic compounds were recovered) and therefore less efficient for recovering of the whole aroma profile, than the traditional solvent extraction. The quantitative determination of DMHF and mesifuran in fresh and processed fruits can also be carried out using a stable isotope dilution assay (Sen, Schieberle, & Grosch, 1991), which provides a high degree of sensitivity and accuracy. In a comparison of the industrial recovery of strawberry flavour by fractional distillation and plate condensation, 38 volatile indicator compounds were identified (Kollmannsberger & Berger, 1994) including mesifuran but not DMHF. However conventional fractional distillation gave a higher yield of the sensorially and analytically preferred aroma concentrate (26% as opposed to 15%).

Probably the best method so far reported for the analysis of DMHF, DMHF glucoside and mesifuran in strawberries is that reported by Sanz et al. (1994) where DMHF, DMHF glucoside and mesifuran were determined simultaneously by an HPLC–UV method after clarification of the juice with Celite and Carrez I and II solutions. The great advantage of this method is its simplicity and speed; the whole analysis can be completed in less than 90 min. Different amounts of the three furanones during the ripening of strawberries have been found in seven cultivars (Sanz, Richardson, & Perez, 1995): the relation of the DMHF and mesifuran levels to the overall aroma evaluations and preferences showed that the best correlation values between DMHF content and strawberry aroma were for Parker and Benton strawberry cultivars.

1.2. Furanones induced by exogenous processes

In a study of “sweet” aroma components for three types of jam, Sugawara, Ito, and Odagiri (1982) showed that the presence of DMHF in the aroma concentrate of

strawberry jam explains why this jam has a sweeter odour than grape or blueberry jam. Given that DMHF is present in the starting slurries of all these fruits, the sweeter odour of strawberry jam could be explained by the major contribution of Maillard reactions that most probably take place during the distillation of the fruit slurries. This proposal was supported by the fact that the aroma concentrate of each jam has a sweeter aroma than that of the corresponding slurry in organoleptic tests.

Several heterocyclic compounds, including DMHF, have been identified in cognac where the amount increased with the age of the brandy (Pisarnitskii, Egorov, Egofarova, & Erygin, 1979). The concentrations of the heterocyclic compounds were correlated with the concentrations of ascorbic acid, which was proposed as the most probable precursor of carbocyclic and heterocyclic compounds accumulating during the ageing of brandy. Schieberle (1992) demonstrated the formation of DMHF from sugar phosphates by heating 0.2 mmol buffered solutions of sugar phosphates at 100 or 150°C for 60 or 45 min, respectively and proposed a possible role for ascorbic acid by showing that addition of ascorbic acid enhanced by a factor of 2 the amounts of DMHF formed from fructose-1,6-diphosphate and fructose-6-phosphate.

DMHF has been claimed to be a predominant odour compound in pastry made with margarine as determined by comparison of flavour profiles of puff-pastries using aroma extraction dilution analysis (AEDA) (Gas-senmeier & Schieberle, 1994). Numerous heterocyclic flavour compounds are formed during roasting of raw foods and many investigations have been devoted to gaining an understanding of this process (Cerny & Grosch, 1992; Semmelroch, Laskawy, Blank, & Grosch, 1995; Ziegler, 1991). Thirteen compounds, including DMHF, were identified by GC–MS in an aroma concentrate prepared from sponge cake by Takei (1977) who used several model systems to study the conditions necessary to produce DMHF; different combinations of glucose, arginine, egg lecithin and sodium carbonate were tested. Roasted almond volatiles were separated on the basis of carbonyl and non-carbonyl fractions and each fraction was analysed by GC–MS (Takei & Yamanishi, 1974). Twenty five compounds were identified and DMHF seemed to make the largest contribution to the sweet aroma of roasted almond. In an analogous study of the flavour extracts of raw and roasted cocoas by GC–MS, 20 new flavour compounds were identified for the first time as cocoa volatiles, among them DMHF and several other heterocyclic compounds (Ziegler, 1991). Also, sugar degradation products, DMHF included, which are formed during drying and roasting of cocoa, important for the flavour development, are discussed. In similar studies with different varieties of roasted coffees, the profiles of volatile compounds were determined and compared for

Robusta and Arabica type coffees (Semmelroch et al., 1995; Tressl, Bahri, Koeppler, & Jensen, 1978). Tressl et al. found 11 diphenols and 7 caramel compounds which were extracted from roasted coffee with pentane–ether, whereas Semmelroch et al. determined 14 important aroma compounds in roasted coffee. Both studies reported higher DMHF levels in Arabica type coffee.

Cerny and Grosch (1993) have studied the effect of the material of cooking utensils on the profile of the flavour formed during roasting. They compared beef flavour roasted in glass to that cooked in stainless steel. The latter material resulted in a higher cooking temperature which led to the formation of a more caramellized flavour. Further studies on the aroma of processed beef by AEDA have shown that DMHF as well as two pyrazines and a thiazoline are particularly important compounds for the roasted, caramel-like and earthy odour notes of roasted and stewed beef (Cerny & Grosch, 1992, 1993; Guth & Grosch, 1994).

Kallio (1988) compared the levels of DMHF in two maple syrups and one birch syrup and concluded that the DMHF concentration was higher in the birch syrup. The difference between the two types of syrups may be related to the larger amounts of fructose and glucose in the birch syrup. Kallio, Rine, Pangborn, and Jennings (1987) have demonstrated that heating birch syrup caused the reduction of the aroma intensity of vanillin but had little effect on the amounts of DMHF and increased the development of burnt aromas. Kallio, Leino, and Salorinne (1989) also reported that the identification of DMHF in the headspace of birch syrup was possible by using headspace GC–MS analysis; DMHF was also found to be one of the major flavour components of Finnish birch syrup (0.8–2.4 mg/kg) (Kallio, 1989).

For orange juice, storage at 35°C for 12 weeks resulted in the formation of DMHF above its taste threshold level (Tatum, Nagy, & Berry, 1975). Walsh, Rouseff, Naim, and Zehavi (1995) have shown that the addition of L-cysteine to the juice reduced the levels of DMHF formed at all temperatures and time-periods tested. Naim et al. (1993) found that small amounts (25 mM) of L-cysteine and N-acetyl-L-cysteine reduced browning and DMHF formation in orange juice stored at 45°C for up to 14 days due to inhibition of Maillard-type reactions by thiols. In the case of aseptically concentrated orange juice, processing at three holding temperatures (84, 87 and 90°C) for 72 s did not affect the amount of DMHF (Rassis and Saguy, 1995). However, DMHF concentration decreased during the storage of the juice at 32°C for 7 weeks.

1.3. Studies on the thermally induced formation of DMHF

For the synthesis of DMHF (Mazenod, Dealy, & Naf, 1992; Wong, Mazenod, & Whitesides, 1983), 6-deoxy-

ketohehexoses (6-deoxy-D-fructose) and methyl α -D-glucopyranoside, respectively, were used. Given the potential of the Maillard reaction to generate heterocyclic compounds (including pyrans and furans), it is not surprising that DMHF formation due to heat has also been studied (Baltes, Kunert-Kirchhoff, & Reese, 1989) and a mechanism for formation of oxygen-containing heterocyclic compounds from D-glucose has been suggested by Kroh (1994). The reaction proceeds principally via 1,2- or 2,3-enolisation of the sugar and β -elimination of water leading to the formation of the 1-, 3- and 4-hexosuloses. Subsequent intramolecular cyclisation and further dehydration gives rise to hydroxymethylfurfural, hydroxyacetyl-furan or DMHF (Fig. 2).

An in-depth study of the formation of DMHF in heat-processed foods was reported by Schieberle (1992) who elucidated the effect of the nature of the heated sugar on the amounts of DMHF formed and suggested a possible thermal pathway for the formation of DMHF. The key precursor of DMHF was fructose-1,6-diphosphate (FDP) which gives rise to acetylformoine and subsequent dehydration of the latter to form DMHF. However, similar amounts of the furanone were obtained when fructose-6-phosphate was used and therefore fructose-6-phosphate appears equally important.

Several studies have focused on the reactions between different amino acids and sugars (Silwar, 1992; Kunert-Kirchhoff & Baltes, 1990; Eiserich, Macku, & Shibamoto, 1992). The aroma produced by the reaction between glucose and proline at 200°C has been reported to be due primarily to five compounds: diacetyl, 2-acetyl-1,4,5,6-tetrahydropyridine, 2-acetyl-1-pyrroline, 2-acetyl-3,4,5,6-tetrahydropyridine and DMHF (Roberts & Acree, 1994). In a study of the reaction of cysteine with glucose (Eiserich et al., 1992), the antioxidative activity of microwave-induced volatile compounds was evaluated; DMHF was the strongest antioxidant. In a similar study on the reaction of cysteine with rhamnose (Silwar, 1992), it was found that the model system cysteine/rhamnose roasted at 200–220°C produced a complex mixture of compounds (180 compounds were detected of which DMHF and 5-methyl furfural were judged to be the most important odorants). This study also revealed that, under roasting conditions, the furan oxygen was substituted by sulphur from H₂S released from cysteine and gave rise to thiophenes. The formation of DMHF by the reaction between pentose sugars and amino acids was investigated by Blank and Fay (1996). When either xylose, ribose or arabinose was heated with glycine or L-alanine at 90°C for 1 h, DMHF was formed as well as directly from pentoses. Experiments using ¹³C-labeled glycine and alanine suggest the incorporation of the Strecker degradation product, formaldehyde, into the pentose moiety thus forming DMHF. The proposed mechanism of DMHF formation

amounts in the microwaved samples, although DMHF concentrations were six-fold higher in the conventionally heated sample.

Apart from these studies with model systems, Nagy, Rouseff, and Lee (1989) have studied the effect of thermal processing and prolonged product storage on the formation of DMHF in citrus juice products. The heat-induced changes (100°C for 30 min) for the most odour-active volatiles in strawberries (DMHF included) were reported by Schieberle (1994), who found by AEDA that the concentrations of nine compounds increased so dramatically that they appeared as new odorants in the heated fruit material: of these compounds (E,E)-2,4-decadienal and (E)- β -damascenone were the most odour-active. The results implied that these compounds were liberated from precursors during the heating process. The concentration of DMHF in the fresh and heated strawberry juice was found to be 16.2 and 29.4 (mg/kg), respectively. It was concluded that the decrease in the amounts of the grass-like (Z)-3-hexenal and the fruity-smelling esters, ethyl- and methyl-butanoate, 2-methyl- and 3-methyl-butanoate, in combination with the increase in concentrations of (E)- β -damascenone, (E,E)-2,4-decadienal and the caramel-like smelling DMHF, are the components mainly responsible for the flavour produced during the heating of strawberries.

The stability of both DMHF and mesifuran has been studied in order to assess whether there was any loss of these furanones during processing and canning of fruits (Hirvi, Honkanen, & Pyysalo, 1980). It was found that the decomposition of both furanones followed first-order kinetics and is pH-dependent, with optimum stability at pH 4. Saccharose and ethyl alcohol had no effect on the stability of the furanones over a concentration range (0–20%). In an additional study of the stability of naturally occurring DMHF derivatives (Roscher et al., 1997d), the half lives of all the DMHF derivatives were determined in aqueous buffer solutions at different pH values (from pH 2.0 to 8.0 at intervals of 1.5 pH units). It was found that the greatest stability of all the DMHF derivatives occurred at pH 3.5–5.0 (the natural pH value of strawberries is in the range 3.5–4.0). 2,5-Dimethyl-4-hydroxy-2H-furan-3-one glucoside was stable from pH 2.0–8.0 and might be unaffected in juices or concentrates over a long storage period. Hydrolysis of malonyl-glucoside cannot take place directly by a β -glucosidase but requires an initial malonyl-esterase activity with the subsequent release of free DMHF in a second glycolytic step.

2. The synthesis of 2,5-dimethyl-4-hydroxy-2H-furan-3-one and its precursors

The chemical synthesis of DMHF has been studied because of its high commercial value as a food flavouring.

The first report of a chemo-enzymatic synthesis of 6-deoxy-D-fructose and 6-deoxy-L-sorbose was that of Hough and Jones (1952). Racemic lactaldehyde was reacted with dihydroxyacetone phosphate (DHAP) to give the two methyl pentoses. The chemical synthesis of DMHF from 6-deoxy-ketohexoses starting with threonine and FDP has been reported (Wong et al., 1983). Threonine was reacted with ninhydrin at 100°C to form lactaldehyde (Zagalak, Frey, Karabatsos, & Abeles, 1966) and the enzymes biphosphate aldolase and triose phosphate isomerase were used to liberate DHAP in situ from FDP. The generated DHAP was then allowed to react with lactaldehyde, via an aldol condensation to produce 6-deoxy-D-fructose (in the case of D-lactaldehyde) or 6-deoxy-L-sorbose (in the case of L-lactaldehyde) (Fig. 3). Conversion of these deoxy sugars to DMHF was achieved by heating at 80°C for 20 h with piperidine. The crude yield of DMHF for this reaction was 78% (Wong et al.). However, the disadvantage of this procedure means that it is almost impossible to control the concentration of lactaldehyde (the immediate precursor of the 6-deoxy-hexoses) which is a very reactive compound that readily dimerises (Chrysochoou, 1973).

In order to carry out in vivo metabolic studies of DMHF, ¹⁴C-monolabelled DMHF was synthesized by applying the procedure outlined in the previous paragraph (i.e. reaction of a 6-deoxy hexose with piperidine, Wong et al., 1983) to fucose L-[1-¹⁴C]. The yield of this reaction was 40% and the monolabelled DMHF obtained had a radiochemical purity of 95% which decreased, however, to 30% within 3 days of storage in aqueous solution at –20°C (Roscher, Schreier, & Schwab, 1997b).

Given the significance of 6-deoxy-ketohexoses as DMHF precursors, other studies on their synthesis have been reported (Drueckhammer et al., 1991; Durrwachter, Drueckhammer, Nozaki, Sweers, & Wong, 1986a; Durrwachter, Sweers, Nozaki, & Wong, 1986b; Fessner, Badia, Eyrisch, Schneider, & Sinerius, 1992; Hecquet, Bolte, & Demuynck, 1994; Toone, Simon, Bednowski, & Whitesides, 1989; Wong & Whitesides, 1985) but all have followed the same synthetic approach despite its main deficiency—i.e. the amount of lactaldehyde, prior to its aldol condensation with DHAP, was not measured. Therefore, only the overall yield can be determined and not the yield of each individual step.

A different biochemical synthetic approach has been used for the preparation of 6-deoxy-hexoses, the immediate DMHF precursors; transketolase has been used for synthesising carbohydrates (Kobori, Myles, & Whitesides, 1992) and, in particular, 6-deoxy-D-fructose and 6-deoxy-L-sorbose (Hecquet et al., 1994). In this approach, (R,R) and (R,S)-2,3-dihydroxybutyaldehyde were separately reacted with hydroxypyruvate using spinach leaf transketolase as catalyst to give 6-deoxy-D-fructose

of Fellous and George (1992) who reacted 2,5-dimethyl-3(2*H*)-furanone with hydrogen peroxide in the presence of a base.

3. The biosynthesis of 2,5-dimethyl-4-hydroxy-2*H*-furan-3-one

Linking the presence of DMHF to that of DMHF glucoside could be a useful step in the search for the biosynthetic pathway of DMHF, as it is suggested that the glucoside is the probable precursor of the free aglycone (Zabetakis & Holden, 1996). DMHF is a component of the male sex pheromone of the insect *Eurycotis floridana* (walker), where the male calling behaviour is associated with the sex pheromone released by the anterior part of tergites 2, 7 and 8. Glandular extracts of tergite 7 possess a characteristic odour of caramel, which is attractive at a distance for the females; DMHF and 4-hydroxy-5-methyl-3(2*H*)-furanone have been identified as the compounds responsible for this odour (Farine, Le Quere, Duffy, Semon, & Brossut, 1993). It is suggested that these insects and many plant species synthesize DMHF for their propagation by attracting either insects of the other sex (in the case of walker flies) or insects for pollination of plants.

The biosynthesis of DMHF has been investigated in *Zygosaccharomyces rouxii* and strawberry callus cultures. In the former case, the quantitative effect of different precursor sugars on the yield of DMHF from the yeast cultures was evaluated and it was proposed that 6-deoxyketoses and FDP are the main precursors of DMHF in this system (Hecquet, Sancelme, Bolte, & Demuynck, 1996) but a biosynthetic pathway to DMHF was not proposed.

In an analogous study with strawberry callus cultures by Zabetakis and Holden (1996), it was indicated the intrinsic potential of strawberry cells to produce DMHF glucoside when the cells were provided with a suitable precursor (6-deoxy-D-fructose). This study also showed the possible important role of 6-deoxyketoses in DMHF biosynthesis. Therefore, further investigations were required concerning the factors controlling the bioformation and bioavailability of these deoxy sugars. In such a study, it was revealed that the deoxy-hexoses, 6-deoxy-L-mannose (rhamnose), 6-deoxy-D-galactose (D-fucose) and 6-deoxy-L-galactose (L-fucose) could be probable precursors of the furan ring of DMHF since, when these hexoses were fed to strawberry calli, the production of DMHF-glucoside was observed, by HPLC-PDA. Of the four hexoses tested, 6-deoxy-D-fructose supplementation resulted in the highest amount of DMHF-glucoside and this may be explained on the basis that it is the aldose or ketose nature of the deoxy sugar, and not the configuration (D- or L-) which mainly governs product formation (Zabetakis et al., 1996).

Recently, Roscher et al. (1997a) have claimed using the ratios [^{13}C]/[^{12}C] of rhamnose, that there is a similar distribution of these isotopes in DMHF formed by *Pichia capsulata* and thus rhamnose may be the precursor of DMHF.

Lactaldehyde has been proposed as the precursor of 6-deoxy-fructose-1-phosphate, and, thereafter, DMHF in strawberries (Zabetakis & Holden, 1995) and therefore 1,2-propanediol has been considered as a possible important precursor in a DMHF biosynthetic pathway. 1,2-Propanediol has been identified in strawberry and its role as a flavour precursor discussed (Zabetakis & Gramshaw, 1997); exogenous 1,2-propanediol enhanced the formation of DMHF-glucoside in strawberry callus cultures. As alcohol dehydrogenases occur in strawberries (Mitchell & Jelenkovic, 1995; Yamashita, Iino, & Yoshikawa, 1978), it is possible for 1,2-propanediol to be oxidised to lactaldehyde which could react with DHAP, an omnipresent key metabolic compound, to produce 6-deoxy-fructose-1-phosphate (Zabetakis & Gramshaw, 1997). This deoxy sugar may in turn be converted to DMHF-glucoside, as suggested by Zabetakis and Holden (1995). The final steps of a putative DMHF biosynthetic pathway are outlined in Fig. 4.

Interestingly, when lactaldehyde was supplemented in the culture medium of strawberry calli, free DMHF, but not the glucoside, was formed. It can, thus, be proposed that there may be two, possibly parallel, biosynthetic pathways to this furanone; one leading to the free aglycone and one to DMHF glucoside (Zabetakis, Moutevelis-Minakakis, & Gramshaw, 1998). Given the presence of 1,2-propanediol in strawberry (Zabetakis & Gramshaw, 1997), the effect of *Methylobacterium extorquens*, a bacterial strain with the capability to oxidise primary and secondary alcohols to the corresponding carbonyl compounds, on the DMHF bioformation in strawberry calli was investigated by treating strawberry cell cultures with these bacteria. It was found that this treatment resulted in the formation of DMHF and mesifuran by strawberry callus cultures (Zabetakis, 1997). In this study, it was also found, using HPLC-PDA on the dinitrophenylhydrazones, that when bacterial suspension cultures were supplemented with 1,2-propanediol, lactaldehyde was formed; this microbially-derived aldehyde might then be used by the plant cells for the biosynthesis of the furanones.

Roscher et al. (1997c) have suggested that DMHF is the precursor of DMHF glucoside based on a 0.7% incorporation of ^{14}C -DMHF to ^{14}C -DMHF glucoside. These authors suggested that DMHF is the precursor of DMHF-glucoside and not vice versa, a statement which might be true in the case of strawberry fruit. However, since the glucosidation and the glycolysis reactions are readily reversible, DMHF-glucoside could be the precursor of DMHF in strawberry calli (Zabetakis & Holden, 1996). In strawberry calli, the endogenous




Fig. 4. Proposed outline for the biosynthesis of DMHF (adapted from Zabetakis et al., 1996).

amounts of DMHF and derivatives are zero and the size of the metabolic pool of each furanone should be thus defined only by the metabolism of exogenously supplied precursors. The callus cultures might therefore be a better biological system for the study of glucosidation and glycolysis reactions involving DMHF and derivatives. In view of that, further experiments with strawberry callus cultures and labelled precursors are needed in order to elucidate the mechanism of the formation of DMHF glucoside.

In summary, in this review, the analytical methods used for the analysis of DMHF in various sources have been evaluated. Given that DMHF is thermolabile, HPLC–UV is probably the most appropriate method for reliable quantitative results. The different synthetic

routes to DMHF were also reviewed in order to explore possibilities for the efficient synthesis of isotopically labelled DMHF and potential precursors of it. These labelled compounds should be of great significance in elucidating the chemical/biochemical pathways of DMHF formation in any system where DMHF is a key flavour component. Lastly, the studies on DMHF biogenesis were also critically presented and points of interest, where future work should be focused, have been suggested.

Acknowledgement

I.Z. is grateful to the Hellenic State Scholarships Foundation for the award of a Research Studentship.

References

- Baltes, W., Kunert-Kirchhoff, J., & Reese, G. (1989). Model reactions on generation of thermal aroma compounds. In: T.H. Parliment, R. J. McGorin & C. T. Ho. (Eds.), *Thermal Generation of Aromas* (pp. 143–155.) Washington, DC: ACS.
- Blank, I., & Fay, L. B. (1996). Formation of 4-hydroxy-2,5-dimethyl-3(2H)-furanone and 4-hydroxy-2(or 5)-ethyl-5(or 2)-methyl-3(2H)-furanone through Maillard reaction based on pentose sugars. *Journal of Agricultural Food Chemistry*, 44, 531–536.
- Blank, I., & Schieberle, P. (1993). Analysis of the seasoning-like flavour substances of a commercial lovage extract (*Levisticum officinale* Koch.). *Flav. Fragr. J.*, 8, 191–195.
- Briggs, M. A., Haines, A. H., & Jones, H. F. (1985). Synthesis of 4-hydroxy-2,5-dimethylfuran-3(2H)-one (Furaneol) from (2R,3R)-tartaric acid. *J. Chem. Soc. Perkin Trans. 1*, 1985, 795–798.
- Brueche, G., Schmarr, H. G., Bauer, A., Mosandl, A., Rapp, A., & Engel, L. (1991). Stereoisomeric aroma compounds. LI. Stereo-differentiation of chiral furanones. Possibilities and limits to origin assignment of aroma compounds. *Z. Lebensm. Unters. Forsch.*, 193, 115–118.
- Buchi, G., Demde, E., & Thomas, A. F. (1973). Syntheses of 2,5-dimethyl-4-hydroxy-2,3-dihydrofuran-3-one (furaneol), a flavor principle of pineapple and strawberry. *J. Org. Chem.*, 38, 123–125.
- Buttery, R. G., Takeoka, G. R., Krammer, G. E., & Ling, L. C. (1994). Identification of 2,5-dimethyl-4-hydroxy-3(2H)-furanone (furaneol) and 5-methyl-4-hydroxy-3(2H)-furanone in fresh and processed tomato. *Lebensmittel Wissenschaft und Technologie*, 27, 592–594.
- Buttery, R. G., Takeoka, G. R., & Ling, L. C. (1995). Furaneol: odor threshold and importance to tomato aroma. *Journal of Agricultural Food Chemistry*, 43, 1638–1640.
- Cerny, C., & Grosch, W. (1992). Evaluation of potent odorants in roasted beef by aroma extract dilution analysis. *Z. Lebensm. Unters. Forsch.*, 194, 322–325.
- Cerny, C., & Grosch, W. (1993). Quantification of character-impact odour compounds of roasted beef. *Z. Lebensm. Unters. Forsch.*, 196, 417–422.
- Cesare, L. F.-di, & Nani, R. (1992). Analysis of volatile compounds of “Isabella” (*Vitis vinifera* × *Vitis labrusca*) red grape juice. *Ital. J. Food Sci.*, 4, 279–284.
- Chrysochoou, P. (1973). Study on the stereospecificity of addition reactions to R- α -hydroxypropionaldehyde. PhD thesis, University of Athens.
- Douillard, C., & Guichard, E. (1990). The aroma of strawberry (*Fragaria ananassa*): characterisation of some cultivars and influence of freezing. *Journal of the Science of Food and Agriculture*, 50, 517–531.
- Drueckhammer, D. G., Hennen, W. J., Pederson, R. L., Barbas, C. F., Gautheron, C. M., Krach, T., & Wong, C. H. (1991). Enzyme catalysis in synthetic carbohydrate chemistry. *Synthesis*, 1991, 499–525.
- Durrwachter, J. R., Drueckhammer, D. G., Nozaki, K., Sweers, H. M., & Wong, C. H. (1986a). Enzymatic aldol condensation/isomerization as a route to unusual sugar derivatives. *J. Am. Chem. Soc.*, 108, 7812–7818.
- Durrwachter, J. R., Sweers, H. M., Nozaki, K., & Wong, C. H. (1986b). Enzymatic aldol reaction/isomerization as a route to unusual sugars. *Tetr. Letters*, 27, 1261–1264.
- Eiserich, J. P., Macku, C., & Shibamoto, T. (1992). Volatile antioxidants formed from an L-cysteine D-glucose Maillard model system. *Journal of Agricultural Food Chemistry*, 40, 1982–1988.
- Farine, J. P., Le Quere, J. L., Duffy, J., Semon, E., & Brossut, R. (1993). 4-Hydroxy-5-methyl-3(2H)-furanone and 2,5-dimethyl-4-hydroxy-3(2H)-furanone, two components of the male sex pheromone of *Eurycotis floridana* (Walker) (Insecta, Blattellidae, Poly-zosteriinae). *Biosci. Biotech. Biochem.*, 57, 2026–2030.
- Fehl, A. J., & Marcott, C. (1989). Capillary gas chromatography/Fourier transform infrared spectroscopy using an injector/trap and liquid/liquid extraction. *Analytical Chemistry*, 61, 1596–1598.
- Fellous, R., & George, G. (1992). Method for preparing 2,5-dimethyl-4-hydroxy-3(2H)-furanone. French patent application FR 90-08772 A1.
- Fessner, W. D., Badia, J., Eyrisch, O., Schneider, A., & Sinerius, G. (1992). Enzymatic syntheses of rare ketose 1-phosphates. *Tetr. Letters*, 33, 5231–5234.
- Gassenmeier, K., & Schieberle, P. (1994). Comparison of important odorants in puff-pastries with butter or margarine. *Lebensmittel Wissenschaft und Technologie*, 27, 282–288.
- Guedes de Pinho, P., & Bertrand, A. (1995). Analytical determination of furaneol (2,5-dimethyl-4-hydroxy-3(2H)-furanone). Application to differentiation of white wines from hybrid and various *Vitis vinifera* cultivars. *Am. J. Enol. Vitic.*, 46, 181–186.
- Guth, H., & Grosch, W. (1994). Identification of the character impact odorants of stewed beef juice by instrumental analyses and sensory studies. *Journal of Agricultural Food Chemistry*, 42, 2862–2866.
- Hecquet, L., Bolte, J., & Demuyne, C. (1994). Chemoenzymatic synthesis of 6-deoxy-D-fructose and 6-deoxy-L-sorbose using transketolase. *Tetrahedron*, 50, 8677–8684.
- Hecquet, L., Sancelme, M., Bolte, J., & Demuyne, C. (1996). Biosynthesis of 4-hydroxy-2,5-dimethyl-3(2H)-furanone by *Zygosaccharomyces rouxi*. *Journal of Agricultural Food Chemistry*, 44, 1357–1360.
- Hirvi, T., Honkanen, E., & Pyysalo, T. (1980). Stability of 2,5-dimethyl-4-hydroxy-3(2H)-furanone and 2,5-dimethyl-4-methoxy-3(2H)-furanone in aqueous buffer solutions. *Lebensmittel Wissenschaft und Technologie*, 13, 324–325.
- Hong, Y. C., Huang, L. C., Reineccius, G. A., Harlander, S. K., & Labuza, T. P. (1990). Production of aroma compounds from strawberry cell suspension cultures by addition of precursors. *Plant Cell Tiss. Org. Cult.*, 21, 245–251.
- Honkanen, E., Pyysalo, T., & Hirvi, T. (1980). The aroma of Finnish wild raspberries, *Rubus idaeus*, L. *Z. Lebensm. Untersuch. Forsch.*, 171, 180–182.
- Hough, L., & Jones, J. K. N. (1952). The synthesis of sugars from simpler substances. Part IV. Enzymic synthesis of 6-deoxy-D-fructose and 6-deoxy-L-sorbose. *J. Chem. Soc.*, 1952, 4052–4055.
- Hunter, G. L. K., Busek, W. A., & Radford, T. (1974). Volatile components of canned alphonso mango. *Journal of Food Science*, 39, 90–93.
- Idstein, H., & Schreier, P. (1985). Volatile constituents of Alphonso mango (*Magnifera indica*). *Phytochem.*, 24, 2313–2316.
- Inokuchi, T., Matsumoto, S., Tsuji, M., & Torii, S. (1992). Electrohalogenation of propargyl acetates and amides to form the 1,1-dibromo-2-oxo functionality and a facile synthesis of furaneol. *J. Org. Chem.*, 57, 5023–5027.
- Kallio, H. (1976a). Identification of vacuum steam-distilled aroma compounds in the press juice of arctic bramble, *Rubus arcticus* L. *Journal of Food Science*, 41, 555–562.
- Kallio, H. (1976b). Development of volatile aroma compounds in arctic bramble, *Rubus arcticus* L. *Journal of Food Science*, 41, 563–566.
- Kallio, H. (1988). Comparison and characteristics of aroma compounds from maple and birch syrups. In: G Charalambous (Ed.), *Frontiers of Flavor*, (pp. 241–248.) Amsterdam: Elsevier Science.
- Kallio, H. (1989). Aroma of birch syrup. *Journal of Agricultural and Food Chemistry*, 37, 1367–1371.
- Kallio, H., Leino, M., & Salorinne, L. (1989). Analysis of headspace of foodstuffs near room temperature. *J. High Res. Chrom. Chrom. Comm.*, 12, 174–177.
- Kallio, H., Rine, S., Pangborn, R. M., & Jennings, W. (1987). Effect of heating on the headspace volatiles of Finnish birch syrup. *Food Chemistry*, 24, 287–299.
- Karg, J. E. (1983). Pineapple flavour—history, formulation, prospects. *Riechst. Arom. Kosmet.*, 33, 163–165.
- Kobori, Y., Myles, D. C., & Whitesides, G. M. (1992). Substrate specificity and carbohydrate synthesis using transketolase. *J. Org. Chem.*, 57, 5899–5907.

- Kollmannsberger, H., & Berger, R. G. (1994). Industrial recovery of strawberry flavour — fractional distillation versus plate condensation. *Z. Lebensm. Unters. Forsch.*, *198*, 491–494.
- Krammer, G. E., Takeoka, G. R., & Buttery, R. G. (1994). Isolation and identification of 2,5-dimethyl-4-hydroxy-3(2H)-furanone glucoside from tomatoes. *Journal of Agricultural and Food Chemistry*, *42*, 1595–1597.
- Kroh, L. W. (1994). Caramelisation in food and beverages. *Food Chemistry*, *51*, 373–379.
- Kunert-Kirchhoff, J., & Baltes, W. (1990). Model reactions on roast aroma formation. VIII. Volatile reaction products from the reaction of phenylalanine with 2,5-dimethyl-4-hydroxy-3(2H)-furanone (furanol) by cooking in a laboratory autoclave. *Z. Lebensm. Unters. Forsch.*, *190*, 14–16.
- Larsen, M., & Poll, L. (1992). Odour thresholds of some important aroma compounds in strawberries. *Z. Lebensm. Unters. Forsch.*, *195*, 120–123.
- Larsen, M., Poll, L., & Olsen, C.E. (1992). Evaluation of the aroma composition of some strawberry (*Fragaria ananassa* Duch) cultivars by use of odour threshold values. *Z. Lebensm. Unters. Forsch.*, *195*, 536–539.
- Latrasse, A. (1991). Fruits III. In H. Maarse (Ed.), *Volatile compounds in Foods and Beverages* (pp. 329–387). New York: Marcel Dekker.
- Lee, H. S., & Nagy, S. (1987). High performance liquid chromatography of 2,5-dimethyl-4-hydroxy-3(2H)-furanone in pineapple and grapefruit juices. *Journal of Food Science*, *52*, 163–165.
- Mayerl, F., Naf, R., & Thomas, A. F. (1989). 2,5-dimethyl-4-hydroxy-3(2H)-furanone glucoside: isolation from strawberries and synthesis. *Phytochem.*, *28*, 631–633.
- Mazenod, F., Dealy, F., & Naf, F. (1992). 2,5-dimethyl-4-hydroxy-3(2H)-furanone (furanol) from methyl- β -D-glucopyranoside. *Chimia*, *46*, 403–405.
- Miller, P. H., Libbey, L. M., & Yang, H. H. (1973). Loganberry flavor components of commercial essence. *Journal of Agricultural and Food Chemistry*, *21*, 508.
- Mitchell, W. C., & Jelenkovic, G. (1995). Characterizing NAD-dependent and NADP-dependent alcohol dehydrogenase enzymes of strawberries. *J. Amer. Soc. Hortic. Sci.*, *120*, 798–801.
- Mosandl, A., Bruche, G., Askari, C., & Schmarr, H. G. (1990). Stereoisomeric flavor compounds XLIV: enantioselective analysis of some important flavor molecules. *J. High Res. Chrom.*, *13*, 660–662.
- Nagy, S., Rouseff, R. L., & Lee, H. S. (1989). Thermally degraded flavors in citrus juice products. In T. H. Parliment, R. J. McGorin, & C. T. Ho (Eds.), *Thermal Generation of Aromas*, (pp. 331–345.) Washington, DC: ACS.
- Naim, M., Wainish, S., Zehavi, U., Peleg, H., Rouseff, R., & Nagy, S. (1993). Inhibition by thiol compounds of off-flavor formation in stored orange juice. 1. Effect of L-cysteine and N-acetyl-L-cysteine on 2,5-dimethyl-4-hydroxy-3(2H)-furanone formation. *Journal of Agricultural Food Chemistry*, *41*, 1355–1358.
- Pickenhagen, W., Velluz, A., Passerat, J. P., & Ohloff, G. (1981). Estimation of 2,5-dimethyl-4-hydroxy-3(2H)-furanone (furanol) in cultivated and wild strawberries, pineapples and mangoes. *Journal of the Science of Food and Agriculture*, *32*, 1132–1134.
- Pisarnitskii, A. F., Egorov, I. A., Egozarova, R. K., & Erygin, G. D. (1979). Carbohydrate degradation products as aroma-forming substances in cognac. *Appl. Biochem. Microbiol.*, *15*, 673–678.
- Polesello, S., Lovati, F., Rizzolo, A., & Rovida, C. (1993). Supercritical fluid extraction as a preparative tool for strawberry aroma analysis. *J. High Res. Chrom.*, *16*, 555–559.
- Pyysalo, T., Suihko, M., & Honkanen, E. (1977). Odour thresholds of the major volatiles identified in cloudberry (*Rubus chamaemorus* L.) and arctic bramble (*Rubus arcticus* L.). *Lebensmittel-Wissenschaft und Technologie*, *10*, 36–39.
- Rapp, A., Knipser, W., & Engel, L. (1980). A typical aroma compounds in grapes and wines from interspecies hybrid vines. I. Strawberry note. *Vitis*, *19*, 13–23.
- Rapp, A., Knipser, W., Engel, L., & Hastrich, H. (1983). Capillary chromatographic investigations on various grape varieties. In G. Charalambous & G. Inglett (Eds.), *Instrumental Analysis of Food* (pp. 435–454.) Orlando: Academic Press.
- Rassis, D., & Saguy, I. S. (1995). Kinetics of aseptic concentrated orange juice quality changes during commercial processing and storage. *Int. J. Food Sci. Technol.*, *30*, 191–198.
- Re, L., Mauren, B., & Ohloff, G. (1973). Ein einfacher zugang 4-hydroxy-2,5-dimethyl-3(2H)-furanon (furanol), einem aromabestandteil von ananas und erdbeere. *Helv. Chim. Acta*, *56*, 1882–1894.
- Roberts, D. D. & Acree, T. E. (1994). Gas chromatography-olfactometry of glucose-proline Maillard reaction products. In T. H. Parliment, M. J. Morello, & R. J. McGorin (Eds.), *Thermally Generated Flavors* (pp. 71–79.) Washington, DC: ACS.
- Rodin, J. O., Himel, C. M., Silverstein, R. M., Leeper, R. W., & Gortner, W. A. (1965). Volatile flavor and aroma components of pineapple, I. Isolation and tentative identification of 2,5-dimethyl-4-hydroxy-3(2H)-furanone. *Journal of Food Science*, *30*, 280–285.
- Roscher, R., Herderich, M., Steffen, J. P., Schreier, P., & Schwab, W. (1996a). 2,5-Dimethyl-4-hydroxy-3[2H]-furanone 6'-O-malonyl- β -D-glucopyranoside in strawberry fruits. *Phytochem.*, *43*, 155–159.
- Roscher, R., Hilkert, A., Gessner, M., Schindler, E., Schreier, P., & Schwab, W. (1997a). L-Rhamnose: progenitor of 2,5-dimethyl-4-hydroxy-3[2H]-furanone formation by *Pichia capsulata*?. *Z. Lebensm. Unters. Forsch.*, *204*, 198–201.
- Roscher, R., Schreier, P., & Schwab, W. (1997b). A facile synthesis of 2,5-dimethyl-4-hydroxy-3(2H)-furanone [2- (or 5-)methyl 14 C] (Furanol [2-(or 5-) methyl 14 C]). *J. Label. Compds. Radiopharm.*, *39*, 493–499.
- Roscher, R., Schreier, P., & Schwab, W. (1997c). Metabolism of 2,5-dimethyl-4-hydroxy-3[2H]-furanone in detached ripening strawberry fruits. *Journal of Agricultural and Food Chemistry*, *45*, 3202–3205.
- Roscher, R., Schwab, W., & Schreier, P. (1997d). Stability of naturally occurring of 2,5-dimethyl-4-hydroxy-3[2H]-furanone derivatives. *Z. Lebensm. Unters. Forsch.*, *204*, 438–441.
- Roscher, R., Steffen, J. P., Herderich, M., Schwab, W., & Schreier, P. (1996b). Synthesis of phenyl 6'-O-malonyl- β -D-glucopyranoside. Facile preparation of malonylated glycoconjugates. *Journal of Agricultural Food Chemistry*, *44*, 1626–1629.
- Sanz, C., Perez, A. G., & Richardson, D. G. (1994). Simultaneous HPLC determination of 2,5-dimethyl-4-hydroxy-3(2H)-furanone and related flavor compounds in strawberries. *Journal of Food Science*, *59*, 139–141.
- Sanz, C., Richardson, D. G., & Perez, A. G. (1995). 2,5-dimethyl-4-hydroxy-3(2H)-furanone and derivatives in strawberries during ripening. In R. L. Rouseff (Ed.), *Fruit Flavors* (pp. 268–275.) Washington, DC: ACS.
- Schieberle, P. (1992). Formation of furaneol in heat-processed foods. In R. Teranishi, G. R. Takeoka, & M. Guntert (Eds.), *Flavor Precursors* (pp. 164–174.) Washington, DC: ACS.
- Schieberle, P. (1994). Heat-induced changes in the most odour-active volatiles of strawberries. In H. Maarse, & D. G. van der Heij (Eds.), *Trends in Flavour Research* (pp. 345–351.) Amsterdam: Elsevier Science.
- Schreier, P., & Drawert, F. (1981). Aroma composition of fruit wines. I. Strawberry wine. *Chem. Mikrobiol. Technol. Lebensm.*, *7*, 23–27.
- Schreier, P., & Paroschy, J. H. (1981). Volatile constituents from Concord, Niagara (*Vitis labrusca*, L.) and Elvira (*V. labrusca*, L. x *V. riparia*, M.) grapes. *Can. Inst. Food Sci. Technol. J.*, *14*, 112–118.
- Semmelroch, P., Laskawy, G., Blank, I., & Grosch, W. (1995). Determination of potent odourants in roasted coffee by stable isotope dilution assays. *Flav. Fragr. J.*, *10*, 1–7.
- Sen, A., Schieberle, P., & Grosch, W. (1991). Quantitative determination of 2,5-dimethyl-4-hydroxy-3(2H)-furanone and its methyl ether using a stable isotope dilution assay. *Lebensm. Wiss. Technol.*, *24*, 364–369.

- Shaw, P. E., & Berry, R. (1977). Hexose-amino acid degradation studies involving formation of pyrroles, furans and other low molecular weight products. *Journal of Agricultural and Food Chemistry*, 25, 641–644.
- Shaw, P. E., Tatum, J. H., & Berry, R. E. (1969). Base-catalyzed sucrose degradation studies. *Journal of Agricultural and Food Chemistry*, 17, 907–911.
- Shibamoto, T., & Yeo, H. (1994). Flavor in the cysteine-glucose model system prepared in microwave and conventional ovens. In T. H. Parliment, M. J. Morello, & R. J. McGorin (Eds.), *Thermally Generated Flavors* (pp. 457–465.) Washington, DC: ACS.
- Shu, C. K., & Ho, C. T. (1989). Parameter effects on the thermal reaction of cystine and 2,5-dimethyl-4-hydroxy-3(2H)-furanone. In T. H. Parliment, R. J. McGorin, & C. T. Ho (Eds.), *Thermal Generation of Aromas* (pp. 229–241.) Washington, DC: ACS.
- Shu, C. K., Mookherjee, B. D., & Ho, C. T. (1985). Volatile components of the thermal degradation of 2,5-dimethyl-4-hydroxy-3(2H)-furanone. *Journal of Agricultural and Food Chemistry*, 33, 446–448.
- Silwar, R. (1992). Gas-chromatography mass-spectrometry investigation of aroma compounds from the reaction of cysteine and rhamnose under roasting conditions. *Z. Lebensm. Unters. Forsch.*, 195, 112–119.
- Sugawara, E., Ito, T., & Odagiri, S. (1982). Sweet aroma components in three kinds of jam. *J. Agric. Chem. Soc. Japan*, 56, 101–108.
- Takei, Y. (1977). Flavor components of sponge cake and production of 2,5-dimethyl-4-hydroxy-3(2H)-furanone in several model systems. *Agric. Biol. Chem.*, 41, 2361–2368.
- Takei, Y., & Yamanishi, T. (1974). Flavor components of roasted almond. *Agric. Biol. Chem.*, 38, 2329–2336.
- Tatum, J. H., Nagy, S., & Berry, R. E. (1975). Degradation products formed in canned single-strength orange juice during storage. *Journal of Food Science*, 40, 707–709.
- Tonsbeek, C. H. T., Plancken, A. J., & Weerdhof, T. (1968). Components contributing to beef flavor. Isolation of hydroxy-5-methyl-3(2H)-furanone and its 2,5-dimethyl homolog from beef broth. *Journal of Agricultural and Food Chemistry*, 16, 1016–1020.
- Toone, E. J., Simon, E. S., Bednowski, M. D., & Whitesides, G. M. (1989). Enzyme-catalyzed synthesis of carbohydrates. *Tetrahedron*, 45, 5365–5422.
- Tressl, R., Bahri, D., Koeppler, H., & Jensen, A. (1978). Diphenols and caramel compounds in roasted coffees of different varieties. *Z. Lebensm. Unters. Forsch.*, 167, 111–114.
- Walsh, M., Rouseff, R., Naim, M., & Zehavi, U. (1995). Improved chromatographic determination of furaneol and p-vinylguaiacol storage off-flavours in canned orange juice: inhibition by L-cysteine and N-acetyl-L-cysteine. IFT Annual Meeting (p. 74) *FSTA* 27, 10A6.
- Willhalm, B., Stoll, M., & Thomas, A. F. (1965). 2,5-dimethyl-4-hydroxy-2,3-dihydrofuran-3-one. *Chem. Ind. (London)*, 18, 1629–1630.
- Williams, A. A., & Mottram, D. S. (1981). Gas chromatographic analysis of furaneol. *J. High Res. Chrom. Chrom. Comm.*, 4, 421–422.
- Wilson, C. W. III, Shaw, P. E., & Knight, R. J. Jr. (1990). Importance of some lactones and 2,5-dimethyl-4-hydroxy-3(2H)-furanone to mango (*Mangifera indica* L.) aroma. *Journal of Agricultural and Food Chemistry*, 38, 1556–1559.
- Wintoch, H., Krammer, G., & Schreier, P. (1991). Glycosidically bound aroma compounds from two strawberry fruit species, *Fragaria vesca* f. *semperflorens* and *Fragaria* × *ananassa*, cv. Korona. *Flav. Fragr. J.*, 6, 209–215.
- Withopf, B., Richling E., Roscher, R., Schwab W., & Schreier P., (1997). Sensitive and selective screening for 6'-O-malonylated glucoconjugates in plants. *Journal of Agricultural and Food Chemistry*, 45, 907–911.
- Wong, K. C., Lim, C. L., & Wong, L. L. (1992). Volatile flavour constituents of chempedak (*Artocarpus polyphema* Pers.) fruit and jackfruit (*Artocarpus heterophyllus* Lam.) from Malaysia. *Flav. Fragr. J.*, 7, 307–311.
- Wong, C. H., Mazenod, F. P., & Whitesides, G. M. (1983). Chemical and enzymatic syntheses of 6-deoxy-hexoses. Conversion to 2,5-dimethyl-4-hydroxy-2,3-dihydrofuran-3-one (furaneol) and analogues. *J. Org. Chem.*, 48, 3493–3497.
- Wong, C. H., & Whitesides, G. M. (1985). Enzymes as catalysts in synthetic organic chemistry. *Angew. Chem. Int. Ed. Engl.*, 24, 617–718.
- Wu, P., Kuo, M. C., Hartman, T. G., Rosen, R. T., & Ho, C. T. (1991). Free and glycosidically bound aroma compounds in pineapple (*Ananas comosus* L. Merr.). *Journal of Agricultural and Food Chemistry*, 39, 170–172.
- Wu, P., Kuo, M. C., Zhang, K. Q., Hartman, T. G., Rosen, R. T., & Ho, C. T. (1990). Analysis of glycosidically bound 2,5-dimethyl-4-hydroxy-3(2H)-furanone. *Perf. Flav.*, 15, 51–53.
- Yamashita, I., Iino, K., & Yoshikawa S., (1978). Alcohol dehydrogenase from strawberry seeds. *Agric. Biol. Chem.*, 42, 1125–1132.
- Yaylayan, V. A., Forage, N. G., & Mandeville, S. (1994). Microwave and thermally induced Maillard reactions. In T. H. Parliment, M. J. Morello, & R. J. McGorin (Eds.), *Thermally Generated Flavors* (pp. 449–456.) Washington, DC: ACS.
- Zabetakis, I. (1997). Enhancement of flavour biosynthesis from strawberry (*Fragaria* × *ananassa*) callus cultures by *Methylobacterium* species. *Plant Cell Tiss. Org. Cult.*, 50, 179–183.
- Zabetakis, I., & Gramshaw, J.W. (1997). 1,2-Propanediol in strawberries and its role as a flavour precursor. *Food Chemistry*, 61, 351–354.
- Zabetakis, I., Gramshaw, J. W., & Robinson, D. S. (1996). The biosynthesis of 2,5-dimethyl-4-hydroxy-2H-furan-3-one and its derivatives in strawberry. In D. S. Mottram & A. J. Taylor (Eds.), *Flavour Science* (pp. 90–93.) London: RSC.
- Zabetakis, I., & Holden, M. A. (1995). A study of strawberry flavour biosynthesis. In P. Etievant & P. Schreier (Eds.), *Bioflavour 95* (pp. 211–216.) Paris: INRA Editions.
- Zabetakis, I., & Holden, M. A., (1996). The effect of 6-deoxy-D-fructose on flavour bioformation from strawberry (*Fragaria* × *ananassa*, cv. Elsanta) callus cultures. *Plant Cell, Tiss. Org. Cult.*, 45, 25–29.
- Zabetakis, I., & Holden, M. A. (1997). Strawberry flavour: analysis and biosynthesis. *J. Sci. Food Agric.*, 74, 421–434.
- Zabetakis, I., Moutevelis-Minakakis, P., & Gramshaw, J. W. (1999). The role of 2-hydroxypropanal in the biosynthesis of 2,5-dimethyl-4-hydroxy-2H-furan-3-one in strawberry (*Fragaria* × *ananassa*, cv. Elsanta) callus cultures. *Food Chem.*, 64(3), 311–314.
- Zagalak, B., Frey, P. A., Karabatsos, G. L., & Abeles, R. H. (1966). The stereochemistry of the conversion of D and L 1,2-propanediols to propionaldehyde. *J. Biol. Chem.*, 241, 3028–3035.
- Ziegler, G. (1991). Composition of flavor extracts of raw and roasted cocoas. *Z. Lebensm. Unters. Forsch.*, 192, 521–525.